

AIMS Genetics, 3(3): 167-176. DOI: 10.3934/genet.2016.3.167 Received: 14 April 2016 Accepted: 31 August 2016 Published: 02 September 2016

http://www.aimspress.com/journal/Genetics

**Research** article

# Vitamin D receptor gene polymorphisms in Sudanese children with

## type 1 diabetes

#### Khalid Eltahir Khalid\*

Department of Basic Medical Sciences, Faculty of Applied Medical Sciences, Albaha University, P.O.Box: 1988, Saudi Arabia.

\* Correspondence: Email: khatahir@bu.edu.sa; Tel: +0096-650-638-4596.

**Abstract:** Type 1 diabetes mellitus (T1DM) is a T cell mediated autoimmune disease. Vitamin D was found to suppress the incidence of diabetes when bind to its receptor (VDR), probably by suppressing T cell activations. Thus the VDR gene polymorphism may have an impact on pathophysiology of this disease. Since there was no consistent association between VDR polymorphisms and the risk of T1DM, this study aimed to investigate a VDR gene polymorphism in Sudanese children with T1DM. We examined the VDR gene *Bsm*1 (rs1544410), *Apa*1 (rs7975232), and *Taq*1 (rs731236) single nucleotide polymorphisms in 174 children with T1DM, and 56 children as control, and the association of these polymorphisms with the diabetic control. Among study patients, the majority (85.63%) of diabetic patients reported metabolically poor controlled (HbA1c > 8%). As compared with the control, patients with T1DM presented more commonly with *Bsm*1 B allele (p = 0.001; OR 0.283; 95% CI 0.131–0.609) and *Taq*1 T allele (p = 0.05; OR 2.429; 95% CI 1.073–5.496). *Apa*1 A allele was less common in patients with T1DM without statistical difference (p = 0.862; OR 1.085; 95% CI 0.546–2.156). Our study suggests that, *Bsm*1 and *Taq*1 polymorphisms of the VDR gene associated with the prevalence of T1DM.

Keywords: Type 1 diabetes mellitus; Vitamin D receptor; Gene polymorphisms; Sudanese children

#### 1. Introduction

Type 1 diabetes mellitus (T1DM) is a multi-factorial autoimmune disorder, characterized by the destruction of pancreatic beta cells resulting in the deficiency of insulin secretion [1], caused by complex interaction between genetic and environmental factors, accounting for 5–10% of diabetes

cases worldwide [2], and it is common among children and young adults and usually developed early under age 30 years.

It is known that the active vitamin D (1,25-dihydroy vitamin D3), play an important role in the regulation of immune cells proliferation and differentiations, lymphocytes activation, cytokines production [3], and insulin secretion [4]. Vitamin D bioactivity is through the vitamin D receptor (VDR), which is belonging to the nuclear hormone receptors superfamily [5].

The VDR gene is located on the long arm of chromosome 12 (12q12) and 14 (q14), and four common single nucleotide polymorphisms (SNPs) have been identified [6-9] namely *Bsm*1, *Fok*1, *Apa*1 and *Taq*1 polymorphisms (designated as rs1544410, rs10735810, rs7975232, and rs731236 SNP, respectively). VDR polymorphism was found to be associated with many autoimmune disorders in human, such as celiac disease [10], multiple sclerosis [11,12], systemic lupus erythematosus [13-15], rheumatoid arthritis [16,17], and hashimoto,s throiditis [18]. The role of VDR polymorphisms in the pathogenesis of T1DM was not clear, large number of studies [5,7,19-25] has been investigated the association between the aforementioned SNPs and the risk of T1DM but the results were inconsistent.

This study was conducted to investigate the association between VDR polymorphisms (at position *Bsm*1, *Taq*1 and *Apa*1) and susceptibility to T1DM in Sudanese population.

#### 2. Materials and Methods

#### 2.1. Patients and controls

The study encompasses 174 T1DM patients [mean onset age  $7.68 \pm 3.59$  (range, 1–16) years old; 90 female and 84 male] refer to the diabetic clinic at Wad Medani Pediatric hospital which was founded in 1987 to serve children at Gezira state and the out skirt provinces. The patients group was diagnosed according the American Diabetes Association for type 1 diabetes [2]. Fifty six unrelated healthy children [9.50  $\pm$  4.23 (range 4–16) years old; 32 female and 24 male]. The patients were classified into metabolically poor glycemic control with HbA<sub>1c</sub> > 8% and well glycemic control with HbA<sub>1c</sub>  $\leq$  8%. HbA<sub>1c</sub> was assessed by chromatographic-spectrophotometer ion exchange (BioSystems). The demographic and the clinical information were obtained from each subject through well structured questionnaire. Informed consent was obtained from the children guardians or relatives, and the Faculty of Medicine Institutional committee approval was obtained.

#### 2.2. VDR gene analysis

DNA was extracted from the whole blood by using the QIAamp DNA Blood Mini Kit protocol (Qiagen, USA). The target gene of the VDR was extracted from whole gene sequence available at the Pubmed (NCBI Reference Sequence: NG\_008731.1), and amplified by the PCR based on primers specific for *Bsm*1, *Apa*1 and *Taq*1 polymorphisms (designated as rs1544410, rs7975232, and rs731236 SNP, respectively) (Table 1). The PCR condition and digestion used for amplification of *Bsm*1 as follows; 94 °C for 5 min, and 30 cycles using the following temperature profile: 94 °C for 1 min, 56 °C for 1 min, 72 °C for 1 min, and final elongation for 5 min. The PCR products were 825-bp long (B allele), and were digested with Mva12691 (*Bsm*I) at 37 °C for 3 hrs, and then subjected to electrophoresis in 2% agarose gel stained with 5  $\mu$ L ethidium bromide. The lengths of the restriction fragments were 649 and 176 bp (mutant b allele). The PCR condition and digestion

used for amplification of *Apa*1 and *Taq*1 as follows; 94 °C for 10 min, and 30 cycles using the following temperature profile: 94 °C for 1 min, 62 °C for 1 min, 72 °C for 1 min, and final elongation for 5 min. The PCR products were 716-bp long (A or T alleles) and were digested with *Apa*I and *Taq*I at 37 °C and 65 °C for 2 hrs respectively, and then subjected to electrophoresis in 2% agarose gel stained with 5  $\mu$ L ethidium bromide. The lengths of the *Apa*I restriction fragments were 481 and 235 bp (mutant "a" allele), and for *Taq*I were 419 and 297 bp. Five percent of the samples were checked twice for results accuracy.

SNP		PCR primer sequences (5"-3")	Fragment size
	<b>F</b> 1		
Bsm1	Forward	CAA CCA AGA CTA CAA GTA CCG CGT CAG IGA	<i>B</i> : 825 bp
	Reverse	AAC CAG CGG AAG AGG TCA AGG G	<i>b:</i> 649 bp, 176 bp
Apa1	Forward	GGG ACG CTG AGG GAT GGC AGA GC	716 bp
	Reverse	GGA AAG GGG TTA GGT TGG ACA GGA	481 bp, 235 bp
Taq1	Forward	GGG ACG CTG AGG GAT GGC AGA GC	716
	Reverse	GGA AAG GGG TTA GGT TGG ACA GGA	481 bp, 235 bp

Table 1. P	rimers design a	nd DNA fragment	t length of VDR	gene polymorphism.
------------	-----------------	-----------------	-----------------	--------------------

## 2.3. Statistical analysis

A chi-squared test ( $\chi^2$  test) was used to evaluate the associations between different genotypes of *Bsm*1, *Taq*1, and *Apa*1 variants and the disease (T1DM vs controls). The comparisons of the genotype frequencies between groups was analysed using the t-test. Odd ratios (ORs) and 95% confidence intervals (CIs) were calculated for each allele and genotype using logistic regression. The differences were considered significant if the *P*-value less than 0.05.

## 3. Results

The base line characteristics of the study subjects are shown in (Table 2). The male to female ratio in diabetic and control groups was 1:1.1 and 4.3 respectively. Among diabetic patients 59.2% were having family history of diabetes, the mean HbA<sub>1c</sub> among diabetic group was significantly high (P < 0.001) as compared with the control group, on the contrary, BMI which was significantly low among diabetic group, 85.6% of T1DM patients characterized by metabolically poor diabetic control.

Characteristics		Diabetics	Non diabetics
		(N = 174)	(N = 56)
Gender (years)	Male	84	24
	Female	90	32
Age at diagnosis (years)		$11.48 \pm 3.39$	$9.50\pm4.23$
Age at onset of disease		$7.68 \pm 3.59$	-
Duration of diabetes (years)		$3.82 \pm 2.82$	-
Family history of diabetes	Negative	71	40
	Positive	103	16
BMI $(kg/m^2)$		$16.22 \pm 2.37$	$19.10 \pm 4.61*$
$HbA_{1c}(\%)$ (normal rang = 5.4–14.5)		$10.72 \pm 2.18$	$6.05 \pm 1.34*$
Diabetic control	Well (≤8.0%)	25 (14.4%)	-
	Poor (>8.0%)	149 (85.6%)	-

### Table 2. Baseline characteristics of diabetic and control subjects.

\* *p* < 0.001

## 3.1. Genetic set

There was a difference in the VDR genotype and allele frequencies between T1DM patients and the control group. *Bsm*1 *BB* and B allele were significantly (P = 0.004 and p = 0.001 respectively) higher frequency in patients with T1DM (Table 3). The complete cleavage of *Bsm*1 to the target gene into homozygous (wild type BB = 825 bp or mutant bb = 649 bp) (Figure 1). On the other hand, *Taq*1 *TT* genotype (p = 0.012) and *T* allele (p = 0.05) were more frequent among study patients (Table 4), where as its complete cleavage of the target gene into homozygous (wild type *TT* = 712 bp or mutant *tt* = 419 bp) (Figure 2). *Apa*1 AA genotype (p = 0.932) and A allele (p = 0.862) were less frequent in TIDM patients without significant difference (Table 5), its complete cleavage to the target gene generates homozygous (wild type *AA* = 716 bp or mutant aa = 481 bp) (Figure 3). *Bms*1 BB genotype frequency was significantly (p < 0.02) high among patients with poor metabolic control group of T1DM as compared with well metabolic control group (Table 6).

Table 3. The allele frequencies of <i>Bsm</i> 1 in diabetic patients and contro
---

Bsm1	Genotypes			Allele	
Group	Bb	BB	Bb	В	В
Diabetic patients	17 (17%)	51 (51%)	32 (32%)	0.61	0.58
Control	21 (42%)	19 (38%)	10 (20%)	0.42	0.39
<i>P</i> -value	$= 0.004^{a}$			= 0.001	
	$\chi^2 = 11.145$			$\chi^2 = 11.014$	

<sup>a</sup> statistical analysis was performed between BB + Bb and bb; Odd Ratio = 0.283; 95% CI 0.131–0.609.

AIMS Genetics



**Figure 1. Detection of** *Bsm***1 polymorphism by PCR-RFLP method.** Lane 1 and 2: wild-type homozygote (BB), Lane 3 and 4: heterozygote (Bb), Lane 5 and 6: mutant homozygote (bb), and Lane 7: 100-bp DNA marker.



**Figure 2. Detection of** *Taq***1 polymorphism by PCR-RFLP method.** Lane 1 and 2: mutant homozygote (tt), Lane 3 and 4: heterozygote (Tt), Lane 5: wild-type homozygote (TT), and Lane 7: 100-bp DNA marker.



**Figure 3. Detection of** *APa***1 polymorphism using PCR-RFLP method.** Lane 1: 100-bp DNA marker, Lane 2 and 3: mutant homozygote (aa), Lane 4 and 5: heterozygote (Aa), and Lane 6: wild-type homozygote (AA).

Tag1	Genotypes			Allele	
Group	Tt	TT	Tt	Т	Т
Diabetic patients	22 (22%)	63 (63%)	15 (15%)	0.54	0.46
Control	16 (32%)	19 (38%)	15 (30%)	0.41	0.49
<i>P</i> -value	$= 0.012^{a}$			= 0.05	
	$\chi^2 = 8.877$			$\chi^2 = 4.688$	

Table 4. The allele frequencies of *Taq*1 in diabetic patients and controls.

<sup>a</sup> statistical analysis was performed between TT + Tt and tt; Odd Ratio = 2.429; 95% CI 1.073–5.496.

Table 5. The allele frequencies of Apa1 in diabetic patients and controls.

Apa1	Genotypes			Allele	
Group	Aa	AA	Aa	А	А
Diabetic patients	44 (44%)	49 (49%)	7 (7%)	0.68	0.32
Control	21 (42%)	26 (52%)	3 (6%)	0.68	0.32
<i>P</i> -value	$= 0.932^{a}$			= 0.862	
	$\chi^2 = 0.141$			$\chi^2 = 0.054$	4

<sup>a</sup> statistical analysis was performed between AA + Aa and aa; Odd Ratio = 1.085; 95% CI 0.546–2.156.

Genotype frequency		HbA <sub>1c</sub>	HbA <sub>1c</sub> (≤8%)	P value
		(>8%) (n = 87)	group $(n = 13)$	
	BB	55 (63.2 %)	6 (46.2 %)	
Bsm1	Bb	9 (12.3 %)	3 (23.0 %)	0.02 <sup>a</sup>
	Bb	23 (26.4 %)	4 (30.8 %)	
	TT	45 (41.4 %)	7 (53.8%)	
Taq1	Tt	29 (33.3 %)	4 (30.8 %)	0.13 <sup>b</sup>
	Tt	13 (14.9 %)	2 (15.4 %)	
	AA	47 (54.0 %)	7 (53.8 %)	
Apa1	Aa	33 (37.9 %)	5 (38.5 %)	$0.46^{\circ}$
	Aa	7 (8.0 %)	1 (7.7 %)	

Table 6. Frequency of VDR polymorphisms among diabetic control groups.

a,b,c statistical analysis was performed between BB + Bb and bb; TT + Tt and tt; AA + Aa and aa.

#### 4. Discussion

Vitamin D could play a role in the pathogenesis of autoimmune diseases particularly T1DM, with its metabolites could inhibit T cell proliferation and suppress the production of certain pro-inflammatory cytokine such as TNF- $\alpha$  and IL-1 [26-29]. Our study showed high frequency of patients with family history of diabetes. An early family study has supported the association between first degree relatives with T1DM and the risk of developing T1DM [30], while other study indicated that, this association depends on which parent has diabetes [31]. Several studies investigated the association between VDR gene polymorphisms and T1DM in different populations like Greece [5], Taiwanese [9], Germany [1;], Finnish [20], Portuguese [21], Japanese [22], Iranian [23], Dalmatian [24], and Egyptians [25]. Most of the aforementioned studies reported controversial

findings, which were most likely due to ethnic variations in addition to the role of the environmental and genetic factors. Thus far, no study indicated the association between VDR gene polymorphisms and susceptibility to T1DM in Sudanese population. Sudanese regions, particularly central Sudan, is inhabited by ethnically, linguistically and culturally diverse populations which makes the comparison of our results with previous studies from different regions is comparably useful, previous studies combines the genotype data of Sudanese populations with other subordinate countries such as Somalia, Egypt and Uganda [32], in addition to Ethiopia and Fulani [33]. Our study analysed three well characterized VDR polymorphisms (*Bsm*1, *Taq*1, and *Apa*1) among Sudanese T1DM children. We were not being able to investigate the fourth *Fok*1 polymorphism owing to difficulties associated with the enzyme itself.

*Bsm*1 and *Apa*1 are located in the intron between exon 8–9, which seems not affecting VDR protein structure, however, *Bsm*1 may influence VDR mRNA stability because it was found strongly linked with the 3 polyA microsatellite repeat in 3" untranslated region [3,38]. Our results showed both *Bsm*1 *BB* genotype and *B* allele were significantly high in patients than controls and is associated with increased risk of T1DM. This result was consistent with findings from other populations [13,16,35,36]. On the contrary, population study in Iran [14], showed no significant association between *Bsm*1 and susceptibility to T1DM, and was found to have protective role in Greece population [5].

Taq1 is a silent SNP within the 3"- noncoding sequences in exon 9 [37], its exact role in the pathogenesis of T1DM is not well defined. In this study and similar to Bsm1, Taq1 TT genotype and T allele exerted susceptibility action to T1DM, which is agreed with other study populations from Greece and Uruguayan [5,29]. This data was not in agreement with other populations such as Portuguese, Iranian, and Korean, where there was no association found [21,23,39], or there was low frequency in Egyptian population [25]. Our data showed no significant difference in Apa1 between patients and control. In agree with our result, Apa1 genotype and alleles frequencies showed no significant differences between T1DM and the control among Portuguese [21]. In contrary, Apa1 gene polymorphism was found associated with T1DM in Greece and Taiwanese populations [5,7].

#### 5. Conclusion

In conclusion, the VDR *Bsm*1 and *Taq*1 were associated with susceptibility to T1DM among Sudanese population. Our sample size was not large enough to represent the whole population, although it enrols the majority of the patients from central Sudan, nevertheless, the strength of an association between SNPS and T1DM in this study is measured by the odds ratio (OR). Hardy-Weinberg equilibrium for the genotypes was not used in this study because of small sample size and to override violation. Since Sudan is known as small continent and characterized by different ethnic groups, further studies from different regions is necessary to correlate between the genetic and environmental factors, VDR polymorphisms, and T1DM.

#### Acknowledgment

Thanks to Dr. Mohammed Osman Abdelwahid, Department of Molecular Biology, National Cancer Institute, University of Gezira for his contribution.

173

## **Conflict of interest**

The author declares no conflict of interest.

## References

- 1. Barrett JC, Clayton DG, Concannon P, et al. (2009) Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 41: 703-707.
- 2. American Diabetes Association (2005) Diagnosis and classification of diabetes mellitus. *Diabetes Care* 28: S37-S42.
- 3. Uitterlinden AG, Fang Y, Van Meurs JB, et al. (2004) Genetics and biology of vitamin D receptor polymorphisms. *Gene* 33: 8143-8156.
- 4. Wang Q, Xi B, Reilly KH, et al. (2012) Quantitative assessment of the associations between four polymorphisms (*FokI*, *ApaI*, *BsmI*, *TaqI*) of vitamin D receptor gene and risk of diabetes mellitus. *Mol Biol Rep* 39: 9405-9414.
- 5. Panierakis C, Goulielmos G, Mamoulakis D, et al. (2009) Vitamin D receptor gene polymorphisms and susceptibility to type 1 diabetes in Crete Greece. *Clin Immunol* 133: 276-281.
- 6. Capoluongo E, Pitocco D, Concolino P, et al. (2006) Slight association between type 1 diabetes and ""ff<sup>\*\*\*</sup> VDR *FokI* genotype in patients from the Italian Lazio Region. Lack of association with diabetes complications. *Clin Biochem* 39: 888-892.
- 7. Chang TJ, Lei HH, Yeh JI, et al. (2000) Vitamin D receptor gene polymorphism influences susceptibility to type 1 diabetes mellitus in the Taiwanese population. *Clin Endocrinol* 52: 575-580.
- 8. Audí L, Martí G, Esteban C, et al. (2004) VDR gene polymorphism at exon 2 start codon (*FokI*) may have influenced type 1 diabetes mellitus susceptibility in two Spanish populations. *Diabetic Med* 21: 393-394.
- 9. Miyamoto K, Kesterson RA, Yamamoto H, et al. (1997) Structural organization of the human vitamin D receptor chromosomal gene and its promoter. *Mol Endocrinol* 11: 1165-1179.
- 10. San-Pedro JI, Bilbao JR, Perez de Nanclares G, et al. (2005) Heterogeneity of vitamin D receptor gene association with celiac disease and type 1 diabetes mellitus. *Autoimmunity* 38: 439-444.
- 11. Narooie-Nejad M, Moossavi M, Torkamanzehi A, et al. (2015) Vitamin D Receptor Gene Polymorphism and the Risk of Multiple Sclerosis in South Eastern of Iran. *J Mol Neurosci* 56: 572-576.
- 12. Tizaoui K, Kaabachi W, Hamzaoui A, et al. (2015) Association between vitamin D receptor polymorphisms and multiple sclerosis: systematic review and meta-analysis of case-control studies. *Cell Mol Immunol* 12: 243-252.
- 13. Jacobs J, Voskuyl AE, Korswagen LA, et al. (2015) The association between *FOK*-I vitamin D receptor gene polymorphisms and bone mineral density in patients with systemic lupus erythematosus. *Clin Exp Rheumatol* 33: 765.
- 14. Carvalho C, Marinho A, Leal B, et al. (2015) Association between vitamin D receptor (VDR) gene polymorphisms and systemic lupus erythematosus in Portuguese patients. *Lupus* 24: 846-853.
- 15. Xiong J, He Z, Zeng X, et al. (2014) Association of vitamin D receptor gene polymorphisms with systemic lupus erythematosus: a meta-analysis. *Clin Exp Rheumatol* 32:174-181.

174

- 16. Song GG, Bae SC, Lee YH (2016) Vitamin D receptor *FokI*, *BsmI*, and *TaqI* polymorphisms and susceptibility to rheumatoid arthritis : A meta-analysis. *Z Rheumatol* 75: 322-329.
- 17. Tizaoui K, Hamzaoui K (2014) Association between VDR polymorphisms and rheumatoid arthritis disease: Systematic review and updated meta-analysis of case-control studies. *Immunobiology* 220: 807-816.
- 18. Djurovic J, Stojkovic O, Ozdemir O, et al. (2015) Association between *Fok*I, *Apa*I and *Taq*I RFLP polymorphisms in VDR gene and Hashimoto's thyroiditis: preliminary data from female patients in Serbia. *Int J Immunogenet* 42: 190-194.
- 19. Ramos-Lopez E, Brück P, Jansen T, et al. (2007) CYP2R1 (vitamin D 25-hydroxylase) gene is associated with susceptibility to type 1 diabetes and vitamin D levels in German. *Diabetes Metab Res Rev* 23: 631-636.
- 20. Turpeinen H, Hermann R, Vaara S, et al. (2003) Vitamin D receptor polymorphisms: no association with type 1 diabetes in the Finnish population. *Eur J Endocrinol* 149: 591-596.
- 21. Lemos MC, Fagulha A, Coutinho E, et al. (2008) Lack of association of vitamin D receptor gene polymorphisms with susceptibility to type 1 diabetes mellitus in the Portuguese population. *Hum Immunol* 69: 134-138.
- 22. Motohashi Y, Yamada S, Yanagawa T, et al. (2003) Vitamin D receptor gene polymorphism affects onset pattern of type 1 diabetes. *J Clin Endocrinol Metab* 88: 3137-3140.
- Mohammadnejad Z, Ghanbari M, Ganjali R, et al. (2012) Association between vitamin D receptor gene polymorphisms and type 1 diabetes mellitus in Iranian population. *Mol Biol Rep* 39: 831-837.
- 24. Zemunik T, Skrabic V, Boraska V, et al. (2005) *Fok*I polymorphism, vitamin D receptor, and interleukin-1 receptor haplotypes are associated with type 1 diabetes in the Dalmatian population. *J Mol Diagn* 7: 600-604.
- 25. Abd-Allah SH, Pasha HF, Hagrass HA, et al. (2014) Vitamin D status and vitamin D receptor gene polymorphisms and susceptibility to type 1 diabetes in Egyptian children. *Gene* 536: 430-434.
- 26. Chiang JL, Kirkman MS, Laffel LMB, et al. (2014) Type1 Diabetes through the Life Span: A position statement of the American Diabetes Association. *Diabetes Care* 20: 2034-2054.
- 27. Rigby WF, Stacy T, Fanger MW, et al. (1984) Inhibition of T lymphocyte mitogenesis by 1,25-dihydroxyvitamin D3 (calcitriol). *J Clin Investig* 74: 1451-1455.
- 28. Mathieu C, van Etten E, Decallonne B, et al. (2004) Vitamin D and 1,25-dihydroxyvitamin D3 as immunomodulators in the immune system. *J Steroid Biochem Mol Biol* 89: 449-452.
- 29. Muller K, Bendtzen K (1992) Inhibition of human T lymphocyte proliferation and cytokine production by 1,25-dihydroxyvitamin D3. Differential effects on CD45RA<sup>+</sup> and CD45RO<sup>+</sup> cells. *Autoimmunity* 14: 37-43.
- 30. Tattersall RB, Pyke DA (1972) Diabetes in identical twins. Lancet 2: 1120-1125.
- 31. Redondo MJ1, Fain PR, Eisenbarth GS (2001) Genetics of type 1A diabetes. *Recent Prog Horm Res* 56: 69-89.
- 32. Babiker HM, Schlebusch CM, Hassan HY, et al. (2011) Genetic variation and population structure of Sudanese populations as indicated by 15 Identifiler sequence-tagged repeat (STR) loci. *Investig Genet* 2: 12.
- 33. Dobon B, Hassan HY, Laayouni H, et al. (2015) The genetics of East African populations: a Nilo-Saharan component in the African genetic landscape. *Sci Rep* 5: 9996.

- 34. Obi-Tabot ET, Tian XQ, Chen TC, et al. (2000) A human skin equivalent model that mimics the photoproduction of vitamin D3 in human skin. *In Vitro Cell Dev Biol Anim* 36: 201-204.
- 35. Zhang J, Li W, Liu J, et al. (2012) Polymorphisms in the vitamin D receptor gene and type 1 diabetes mellitus risk: an update by meta-analysis. *Mol Cell Endocrinol* 355: 135-142.
- 36. Wang G, Zhang Q, Xu N, et al. (2014) Associations between two polymorphisms (*FokI* and *BsmI*) of vitamin D receptor gene and type 1 diabetes mellitus in Asian population: a meta-analysis. *PLoS One* 9: e89325.
- 37. Morrison NA, Qi JC, Tokita A, et al. (1994) Prediction of bone density from vitamin D receptor alleles. *Nature* 367: 284-287.
- 38. Mimbacas A, Trujillo J, Gascue C, et al. (2007) Prevalence of vitamin D receptor gene polymorphism in a Uruguayan population and its relation to type 1 diabetes mellitus. *Genet Mol Res* 6: 534-542.
- 39. Cheon CK, Nam HK, Lee KH, et al. (2015) Vitamin D receptor gene polymorphisms and type 1 diabetes mellitus in a Korean population. *Pediatr Int* 57: 870-874.



© 2016 Khalid Eltahir Khalid licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0)

AIMS Genetics