

---

*Research article*

## The possible function of Flp1 in homologous recombination repair in *Saccharomyces cerevisiae*

Huong Thi Thu Phung\*, Hoa Luong Hieu Nguyen and Dung Hoang Nguyen

NTT Hi-Tech Institute, Nguyen Tat Thanh University, Ho Chi Minh city, Vietnam

\* **Correspondence:** Email: [ptthuong@ntt.edu.vn](mailto:ptthuong@ntt.edu.vn); Tel: +84981411701.

**Abstract:** *Saccharomyces cerevisiae* Mus81 is a structure-selective endonuclease which constitutes an alternative pathway in parallel with the helicase-topoisomerase Sgs1-Top3-Rmi1 complex to resolve a number of DNA intermediates during DNA replication, repair, and homologous recombination. Previously, it was showed that the N-terminal region of Mus81 was required for its *in vivo* function in a redundant manner with Sgs1; *mus81 $\Delta$ 120N* mutant that lacks the first 120 amino acid residues at the N-terminus exhibited synthetic lethality in combination with the loss of *SGS1*. In this study, the physiologically important role of the N-terminal region of Mus81 in processing toxic intermediates was further investigated. We examined the cellular defect of *sgs1 $\Delta$ mus81 $\Delta$ 100N* cells and observed that although viable, the cells became very sensitive to DNA damaging agents. A single-copy suppressor screening to seek for a factor(s) that could rescue the drug sensitivity of *sgs1 $\Delta$ mus81 $\Delta$ 100N* cells was performed and revealed that Flp1, a site-specific recombinase 1 encoded on the 2-micron plasmid was a suppressor. Moreover, Flp1 overexpression could partially suppress the drug sensitivity of *mus81 $\Delta$*  cells at 37 °C. Our findings suggest a possible function of Flp1 in coordination with Mus81 and Sgs1 to jointly resolve the branched-DNA structures generated in cells attempting to repair DNA damages.

**Keywords:** Flp1; genetic screening; homologous recombination repair; Mus81; Sgs1

---

### 1. Introduction

Mus81, a highly conserved DNA structure-selective endonuclease, is related to the XPF/Rad1 family of proteins involved in DNA nucleotide excision repair. Mus81 functions as a heterodimeric protein complex with a partner, namely Eme1 and Eme2 in humans, Eme1 in fission yeast, and

Mms4 in budding yeast [1–3]. Its partner proteins are indispensable for stability and the nuclease activity of the complex [4].

The *mus81* mutants are hypersensitive to different types of DNA damaging agents including ultra violet irradiation, methyl methanesulfonate (MMS), hydroxide urea (HU), 2-phenyl-3-nitroso-imidazo [1,2- $\alpha$ ] pyrimidine, cisplatin, doxorubicin, tirapazamine, and camptothecin [5–7]. Mus81 can cleave a numerous of branched-DNA structures that may form *in vivo* during many DNA transactions such as nicked Holliday Junctions (HJs), D-loop, replication forks with the lagging strand at the junction point, and 3'-flap [1,3,8]. Despite the fact that Mus81 was reported as a HJs resolvase because the affinity-purified Mus81–Eme1 complex exhibited marked activity on intact HJs [9–11], intact HJs are considered as generally poor substrates for this nuclease [1,3,8,12–14]. Later study proved that the recombinant fission or budding yeast Mus81 complexes exhibited strong activity on intact HJs based on the tetramer formation [15]. In contrast, more recent work demonstrated that *Saccharomyces cerevisiae* Mus81–Mms4 functions as a single heterodimer in recombinational DNA repair and hardly cleaves intact HJs [16]. Interestingly, the activity of Mus81–Mms4 is cell-cycle regulated by the phosphorylation of Mms4 by Cdc28 (CDK) and Cdc5 (Polo-like kinase) which enhance the activity of Mus81 complex on intact HJs [17,18]. Recently, a third cell cycle kinase, Cdc7-Dbf4 (DDK) was shown to be able to phosphorylate Mus81-Mms4 in conjunction with Cdc5 in an interdependent manner and this phosphorylation is also strictly required for Mus81 activation in mitosis [19]. Besides, Mus81-Mms4 was proved to relocalize to subnuclear foci with Rad1-Rad10 and Slx1-Slx4 in response to DNA damage and the relocalization correlated with the function of the complex endonuclease [20].

*S. cerevisiae* *MUS81* and *MMS4* genes were both identified in a synthetic lethality screen of *sgs1* $\Delta$  mutants [21]. Sgs1, a member of the ubiquitous RecQ family of DNA helicases was shown to form a stable complex with Top3 and Rmi1 which enhances the enzymatic activity of Sgs1-Top3 complex. Inactivation of constituents of Sgs1–Top3–Rmi1 complex resulted in the increased level of mitotic recombination, sister chromatid or chromosome exchanges, and genome instability, indicating that the complex plays critical role in homologous recombination repair pathway [22]. Importantly, the synthetic lethality of double deletion of *mus81* or *mms4* together with *sgs1* can be rescued by further deletion of recombination proteins, such as Rad51 or Rad52 [1,8,21,23]. These results prove that Mus81 complex functions downstream of homologous recombination, being significantly involved in processing recombination intermediates in parallel or redundantly with Sgs1 complex [6,8,13,23,24].

Recently, our previous study showed the genetic and functional interaction of Rad27, an important nuclease involving in Okazaki fragment processing and base excision repair, and the Mus81 complex [25,26]. The presence of Rad27 stimulated endonuclease activity of Mus81-Mms4 *in vitro* and vice versa, implicating the joint function of both enzymes in DNA replication or/and recombination. Besides, Srs2, a DNA helicase and DNA-dependent ATPase which is involved in DNA repair and checkpoint recovery, has been currently demonstrated to be able to promote Mus81–Mms4-mediated resolution of recombination intermediates [27]. Also, more recently, several enzymes were shown to be able to enhance the enzymatic activity of Mus81-Mms4 complex, such as heteropentameric replication factor C (RFC) complex and Esc2, an adaptor protein containing no known enzymatic domains, indicating the cooperation between Mus81 and these partners in the process of intermediates [28,29]. The functional interaction of Mus81 complex and its partners depended on their physical interaction, specifically requiring the N-terminal region of Mus81 [26,27].

Moreover, these physical and functional interactions are significantly important for cellular function of Mus81 in both budding yeast [26,27,30] and human [31]. Taking together all, Mus81–Mms4 is most likely to work with other proteins involved in a variety of DNA metabolisms including DNA repair, replication fork stability, and joint molecule formation/resolution during homologous recombination in order to safeguard the genome integrity.

Here, we further investigated the significance role of Mus81 N-terminus *in vivo* by performing a single-copy suppressor screening to seek for a factor(s) that can suppress the cellular defect caused by function loss of Mus81 N-terminal region. Unlike the *mus81<sub>Δ120N</sub>* mutant allele which is synthetic lethality when combined with *sgs1Δ*, the *sgs1Δmus81<sub>Δ100N</sub>* double mutant cells could survive, however, exhibited excessive sensitivity to DNA damaging agents. Through screening, we successfully recovered a candidate that can rescue the HU sensitivity of *sgs1Δmus81<sub>Δ100N</sub>* mutant cells, namely *FLP1*, a site-specific recombinase 1 which is encoded on the 2-micron plasmid. Accordingly, Flp1 is the possible suppressor of cellular defect caused by the dysfunction of Sgs1 and Mus81 when cells are challenged by DNA damages.

## 2. Materials and methods

### 2.1. Yeast strains

*S. cerevisiae* NJY1777 (*MATa ade2-1 ade3::hisG ura3-1 his3-11,15 trp1-1 leu2-3,112 lys2 mus81-10::KAN sgs1-20::hphMX4 can1-100 + pJM500-URA3-SGS1*) was a courtesy from Dr. Miki Ii at University of Alaska Anchorage (AK, USA) [32].

### 2.2. Drop dilution assay

The plasmid pRS325, a yeast episomal vector with a *LEU2* marker, harboring wild-type *MUS81* or *mus81* mutant alleles driven by *ADHI* promoter was transformed into NJY1777. The transformants were grown on plates containing selective synthetic defined media and a single colony from each of the transformants was inoculated into liquid media (1 mL) until saturation. Cell densities were adjusted to  $OD_{600} = 1$  ( $\sim 2 \times 10^7$  cells/mL) by diluting with dH<sub>2</sub>O, followed by spotting of 10-fold serial dilutions onto selective media plates containing with or without DNA damaging agents. The plates were incubated for 4 days at 30 °C. The complementation of *sgs1Δmus81Δ* synthetic lethality by *mus81* mutant alleles was performed in the presence of 5-FOA (uracil analog, 5-fluoroorotic acid). To test the drug sensitivity of *mus81* mutant alleles in the *sgs1Δ* null strain, cells which grew in the presence of 5-FOA were spotted onto plates containing indicated concentrations of DNA damaging agents.

### 2.3. Screening single-copy suppressors of *sgs1Δmus81<sub>Δ100N</sub>* mutant

Yeast genomic DNA library was constructed by Sau3AI-partial digestion of genomic DNA of *S. cerevisiae* YPH499 strain (*MATa ura3-52 lys2-801 ade2-101 trp1-Δ63 his3-Δ200 leu2-Δ1*). The fragmented genomic DNA with estimate 5.6 kb in length on average was ligated into BamHI-digested pRS315 plasmid, a yeast centromere vector with a *LEU2* marker. Ligation product was then transformed into *Escherichia coli* competent cells and the library plasmids were extracted

and stored at  $-80\text{ }^{\circ}\text{C}$  for long-term usage. Before screening, the yeast genomic DNA library was amplified using Plasmid Maxi Kit (QIAGEN, Hilden, Germany).

NJY1777 cell containing a plasmid harboring wild-type *SGS1* gene with a *URA3* marker was transformed with *mus81 $\Delta$ 100N* gene consisted in pRS314 plasmid, a yeast centromere vector with a *TRP1* marker. Transformants were grown in the selective media and transferred onto plates containing 5-FOA, producing double mutant *sgs1 $\Delta$ mus81 $\Delta$ 100N* cells. The double mutant cells were then transformed with yeast genomic DNA library. Transformants were grown in selective media for 24 hours at  $30\text{ }^{\circ}\text{C}$ , followed by replica plating onto the same medium supplemented with 20 mM HU. The plates were incubated for additional 3–4 days. Selected colonies that grew on HU plates were examined for HU resistant capability by drop dilution assay. The HU-resistant colonies were transferred to liquid medium, and total plasmids were isolated. To confirm single-copy suppression, recovered plasmids were retransformed into the *sgs1 $\Delta$ mus81 $\Delta$ 100N* mutant cells and examined for their ability to support cell growth in the presence of HU. Double-checked plasmids were analyzed by sequencing to identify genomic DNA fragments inserted. One of the analyzed plasmids contained the full length of *FLP1* gene.


### 3. Results

#### 3.1. The significance of the N-terminal region of Mus81

We previously found that the *sgs1 $\Delta$ mus81 $\Delta$ 120N* double mutant cell was synthetic lethality similarly to the *sgs1 $\Delta$ mus81 $\Delta$*  double mutant, indicating that the deletion of the first 120 amino acids at the N-terminal region of Mus81 affected the cellular function of this enzyme ([26]; Figure 1A). In order to further define *in vivo* defects associated with the loss of Mus81 N-terminus, we examined the viability of *sgs1 $\Delta$ mus81 $\Delta$ 80N* and *sgs1 $\Delta$ mus81 $\Delta$ 100N* mutant cells using *sgs1 $\Delta$ mus81 $\Delta$*  double mutant cells that were rendered viable by the presence of a plasmid expressing wild-type Sgs1. With the control empty vector transformed, in the presence of 5-FOA the *sgs1 $\Delta$ mus81 $\Delta$*  cells had lost the plasmid containing *SGS1* and became lethal (Figure 1A). Expectedly, these cells developed normally in the presence of episomal copies of wild-type *MUS81* (Figure 1A). In contrast to the synthetic lethality of *sgs1 $\Delta$ mus81 $\Delta$ 120N*, we found that the presence of *mus81 $\Delta$ 80N* and *mus81 $\Delta$ 100N* alleles, despite of poor development, succeeded to support the growth of *sgs1 $\Delta$ mus81 $\Delta$*  cells (Figure 1A). These findings show that a region between the 100<sup>th</sup> and 120<sup>th</sup> amino acid residues at the N-terminus of Mus81 is crucial for the function of Mus81 *in vivo* which is relevant to the cell survival in the absence of Sgs1. Here, it was observed that the deficient growth of the *sgs1 $\Delta$ mus81 $\Delta$ 100N* and *sgs1 $\Delta$ mus81 $\Delta$ 80N* double mutants represented the “bottleneck effect” where a size of population reduced sharply due to a sudden change of environment.

Next, MMS sensitivity of the *mus81 $\Delta$ 80N* and *mus81 $\Delta$ 100N* single mutant cells and the *sgs1 $\Delta$ mus81 $\Delta$ 80N* and *sgs1 $\Delta$ mus81 $\Delta$ 100N* double mutant cells were investigated. When wild-type *MUS81* was introduced into the *mus81 $\Delta$*  deletion cells, they grew normally in the presence of MMS (Figure 1B). Unlike drug sensitivity phenotype of *mus81 $\Delta$*  and *mus81 $\Delta$ 120N* [26], the *mus81 $\Delta$ 80N* and *mus81 $\Delta$ 100N* cells behaved like wild-type (Figure 1B). However, the *sgs1 $\Delta$ mus81 $\Delta$ 80N* and *sgs1 $\Delta$ mus81 $\Delta$ 100N* double mutants were sensitive to MMS and HU treatment (Figure 1C), indicating that the defect causing by the absence of the first 80 and 100 amino acid residues at Mus81 N-terminus, respectively, induced severe troubles to the cell, especially in dealing with DNA-lesion

induced by DNA damaging agents. Overall, these results further demonstrated the critical function of Mus81 N-terminal region in homologous recombination repair pathways when cells are faced with accumulated DNA lesions caused by DNA damaging agents.




**Figure 1.** The cellular defect of *mus81 $_{\Delta 80N}$*  and *mus81 $_{\Delta 100N}$* . (A) The complementation of *sgs1 $\Delta$ mus81 $\Delta$*  synthetic lethality by *mus81* mutant alleles. (B) The MMS sensitivity of *mus81 $_{\Delta 80N}$*  and *mus81 $_{\Delta 100N}$* . (C) The MMS sensitivity of *sgs1 $\Delta$ mus81 $_{\Delta 80N}$*  and *sgs1 $\Delta$ mus81 $_{\Delta 100N}$* . The cells that grew in the presence of 5-FOA (in panel A) were spotted onto without or with indicated amount of MMS or HU concentration.

### 3.2. The single-copy suppressor screening to find out a factor(s) that can suppress the cellular defect causing by the dysfunction of N-terminal region of Mus81

We aim to seek for an alternative pathway that can cope with the loss of function of the important Mus81 N-terminal region. The results in Figure 1A point out that the region between the 100<sup>th</sup> to 120<sup>th</sup> amino acid residues at the N-terminus of Mus81 is indispensable for the cellular function of Mus81 in the absence of Sgs1 because removal of this small portion in the absence of Sgs1 led to cell death unavoidably. To perform the single-copy suppressor screening to define a suppressor of

Mus81 lacking N-terminus mutant, we decided to maintain this small region due to its essential function *in vivo*. Thus, we chose the defective phenotype of *sgs1Δmus81<sub>Δ100N</sub>* cells (Figure 1C) to carry out the suppressor screening in order to identify a factor that can rescue this defect.


As shown in Figure 2, the yeast genomic DNA library was transformed into the *sgs1Δmus81<sub>Δ100N</sub>* cells and the transformants were replicated onto plates containing 20 mM HU. Survival colonies were selected and serial-diluted spotted onto HU-plates in order to evaluate the HU resistant ability. Well growing cells were selected and the transformed plasmids were extracted. The plasmids were next transformed back into the *sgs1Δmus81<sub>Δ100N</sub>* cells and transformants were re-examined for HU-resistant capability. Finally, the plasmids producing resistant transformants were sequenced to define the possible genes involving in suppression of the Mus81<sub>Δ100N</sub> dysfunction.



**Figure 2.** Scheme of the single-copy suppressor screening of drug sensitivity of the *sgs1Δmus81<sub>Δ100N</sub>* cells. (1) transformation with yeast genomic library, (2) replica onto HU plates, (3) and (4) drop dilution assay, (5) sequencing. Abbreviations: P, positive control; N, negative control; C1–4, representative colonies number 1 to 4.

Collectively, after replica plating step, there were fifty-seven colonies that could grow on HU plates. Choosing those colonies and using drop dilution assay, we were able to examine the HU-resistant ability of fifty-four colonies (Figure 3). Among fifty-four checked colonies, forty-one were capable of suppress

HU sensitivity of the *sgs1Δmus81Δ100N* mutant. There were twenty-three strong suppressors in comparison to wild-type cells (Figure 3, Table 1). Next, plasmids from forty-one colonies were extracted and re-transformed into the *sgs1Δmus81Δ100N* cells. Among forty-one candidate plasmids extracted, thirty-eight successfully created transformants. Then transformants were serial-diluted spotted onto plates containing HU to evaluate their survival. Among thirty-eight obtained transformants, only sixteen were capable of resisting to HU treatment (Figure 4, Table 1).



**Figure 3.** A drop dilution assay examining the transformant colonies that survive in the presence of HU. The *sgs1Δmus81Δ100N* cells containing yeast genomic DNA fragment which survived on HU plates after replica step were selected and serial-diluted spotted onto plates without or with 20 mM HU.

**Table 1.** Summary of the single-copy suppressor screening of *sgs1Δmus81Δ100N* mutant.

Candidate	Suppression	Confirmed Suppression	Gene sequence
1	+++	+++	<i>SGS1</i>
2	+	-	
3	+++	+++	<i>SGS1</i>
4	+	-	
5	+	-	
6	++	-	
7	-	/	
8	+	-	
9	NA	/	
10	NA	/	
11	+	++	NA
12	+	-	
13	++	+++	<i>SGS1</i>
14	+	+++	<i>SGS1</i>
15	-	/	
16	NA	/	
17	-	/	
18	+++	+++	<i>SGS1</i>
19	+	-	
20	-	/	
21	+++	+++	<i>SGS1</i>
22	-	/	
23	+++	+++	<i>SGS1</i>
24	+++	+++	<i>SGS1</i>
25	+	+++	<i>SGS1</i>
26	+	-	
27	++	+++	<i>SGS1</i>
28	+	-	
29	+	-	
30	+	-	
31	+	++	<i>FLP1</i>
32	+	-	
33	++++	NA	
34	+++	+++	<i>SGS1</i>
35	+	-	
36	F	/	
37	+++	+++	<i>SGS1</i>
38	+	-	
39	+++	+++	<i>SGS1</i>
40	+++		+++
41	-	/	

*Continued on next page*



Candidate	Suppression	Confirmed Suppression	Gene sequence
42	-	/	
43	-	/	
44	+++	NA	
45	++++	-	
46	++++	/	
47	++	-	
48	++++	-	
49	-	/	
50	+	-	
51	++++	-	
52	+++	-	
53	+++	-	
54	-	/	
55	++++	-	
56	-	/	
57	+	-	
Sum	41	16	

(+) suppressed; (+++) strong suppressed in comparison to positive control; (-) not suppressed.


FFalse positive; <sub>NA</sub>Not available

### 3.3. *Flp1*, a site-specific recombinase 1 encoded on the 2-micron plasmid is a suppressor of *mus81* mutant lacking the N-terminal region

Sixteen plasmids that transformants created were able to grow well in the presence of HU were sequenced to identify the genomic DNA fragments inserted. Finally, it revealed fourteen plasmids harboring *SGS1* sequence including either or both of its upstream and downstream sequence and nearby region on chromosome XIII, jointly forming inserted fragment of approximately 6 kb. The presence of *SGS1* in screening results served as a positive control for our suppressor screening approach. The plasmid number 31 contained an upstream sequence (around 250 base pairs) and full length of a gene called *FLP1*, a site-specific recombinase 1 encoded on the 2-micron plasmid which is a multi-copy selfish extrachromosomal DNA element found in the nucleus in budding yeast. The pRS315 plasmid that we used to construct genomic library is the yeast centromere vector that does not harbor the 2-micron origin of replication and is maintained at a low copy number. Moreover, even yeast episomal plasmids do not contain the full-length Flp1 sequence and its upstream region, indicating that the presence of Flp1 sequence in screening result is definitely not derived from the used plasmid. Here, it is clear that the construct of genomic DNA fragment found in plasmid number 31 could guarantee the expression of functional Flp1 due to the presence of its native promoter in upstream sequence and completed coding sequence.


The 2-micron plasmid is relatively small (approximately 6.3 kb), and resides in most common strains of *S. cerevisiae* at steady-state copy number of 40–60 molecules per haploid cell [33,34]. The plasmid harbors an origin of bidirectional DNA replication called ARS (autonomous replication sequence), *FLP1* gene, three other genes encoding proteins required for regulation of Flp1 expression, namely *REP1*, *REP2*, and *FAF1*, a set of small direct repeats forming a partitioning locus named STB,

and two 599-bp inverted repeat sequences called Flp Recombination Targets (FRTs) [33,35–38]. Flp1 with a couple of FRT sites residing in the plasmid in a head-to-head orientation belong to the amplification system which is responsible for the 2-micron plasmid self-replication [38].




**Figure 4.** A drop dilution assay to confirm the suppression ability of the candidates. The *sgs1Δmus81Δ100N* cells were transformed with extracting plasmids from selected candidates and then serial-diluted spotted onto plates without or with 20 mM HU.

We questioned that the suppression of HU sensitivity by Flp1 overexpression in *sgs1Δmus81Δ* cells expressing Mus81<sub>Δ100</sub> is associated with the loss of N-terminal region of Mus81 which is equivalent to loss of Mus81 function, or the loss of Sgs1, or a combination of these defects. Thus, using the plasmid number 31, we examined the ability of Flp1 overexpression in rescuing cellular defects of the *mus81Δ* and *sgs1Δ* mutants, respectively and found that Flp1 overexpression was unable to suppress the HU sensitivity of *sgs1Δ* (Figure 5). The transformation of plasmid number 31 could not suppress the *mus81Δ* mutant defect at temperature 30 °C, however, it succeeded to make the mutant partially resistant to HU treatment at 37 °C (Figure 5). These findings, together with the screening results, indicate that the suppression by Flp1 is related to the loss of Mus81 function, not Sgs1.



**Figure 5.** Transformation of plasmid number 31 can partially suppress the drug sensitivity of *mus81Δ* at 37 °C. NJY1777 strain was transformed with an empty vector pRS315 or vectors containing *MUS81* or screened plasmid number 31. To create *sgs1Δ* mutant cells, NJY1777 strain was transformed with pRS314 vector containing *MUS81* and transformants were grown in the presence of 5-FOA to remove *pJM500-URA3-SGS1* plasmid.

Next, *FLP1* gene was separately cloned into pRS315 vector and using drop dilution assay, we observed that the Flp1 overexpression could partially rescue the HU sensitivity of the *sgs1Δmus81Δ100N* mutant cells (Figure 6A). This result confirmed that Flp1 is a single-copy suppressor of the *sgs1Δmus81Δ100N* mutant. Besides, overexpression of Flp1 failed to suppress the HU sensitivity of *rad52Δ* and *rad52Δsgs1Δmus81Δ100N* cells (Figure 6B), indicating that Flp1 does not involve in the processing of intermediates at early stage of homologous recombination repair. Also, it was unable to rescue the wild type cells at high amount of HU treatment (Figure 6C), confirming the specific suppressive effect towards *mus81* mutants which lack the full function of Mus81-Mms4 *in vivo*. Flp1 catalyzes a site-specific recombination reaction via interacting with DNA sequences within its FRT sites, producing a single-stranded break on both DNA strands within the target sites that results in inversion of a segment of the 2-micron plasmid [38,39]. Flp1 was determined to remain binding to the 3' side of each break by an O-phosphotyrosyl residue [40]. Accordingly, Flp1 is a very promising suppressor because all of its activities are potentially required in homologous recombination repair pathway when Sgs1 is absent and Mus81 is partially dysfunctional.



**Figure 6.** Suppression of drug sensitivity by Flp1 overexpression is specific for mus81 mutants. (A) Overexpression of Flp1 could partially rescue the HU-sensitivity of the *sgs1Δmus81 $\Delta$ 100N* cells. *FLP1* coding sequence was cloned into pRS315 plasmid and its expression was driven by *ADHI* promoter. Empty vector pRS315 or vector containing *MUS81* or *FLP1* was then transformed into the *sgs1Δmus81 $\Delta$ 100N* cells. (B) Overexpression of Flp1 could not suppress the HU-sensitivity of *rad52Δ* and *rad52Δsgs1Δmus81 $\Delta$ 100N* cells. (C) Overexpression of Flp1 could not rescue the HU-sensitivity of wild-type cells.

#### 4. Conclusions

In mitotic cells, the homologous recombination intermediates are preferentially processed by a ‘dissolution’ activity of the Sgs1-Top3-Rmi1 complex to merely generate non-crossover products. When cells are challenged with DNA damaging agents interfering with normal progression of replication forks or producing excessively double strand breaks, the joint work of Sgs1 and Mus81 could rapidly resolving the excess amount of DNA damages, despite of the risk of crossover product formation. Our previous study has shed light on the important role of Mus81 N-terminal region for

the full function of Mus81 complex in the absence of Sgs1 during homologous recombination repair pathway [26]. Next, we raised a question for the alternative pathway that can rescue the cellular defect of *mus81* mutant isolated in the study. As described in Results section, we have succeeded to recover the *FLP1* gene on the 2-micron plasmid as the suppressor. The 2-micron circle independently reproduces itself with chromosome-like stability via the joined activity of a plasmid amplification system and a plasmid partitioning system. There has not been obvious evidence for the advantage or disadvantage of this plasmid existence to its host [33]. However, a very high copy numbers of the plasmid is harmful to the host, leading to cell cycle misregulation and cell lethality [34]. There is a fact that our strain cells should already harbor 2-micron plasmids but were failed to survive when treated with HU without being further transformed with an extra-plasmid containing *FLP1* sequence. This observation can be explained by regulation of the endogenous 2-micron plasmid number and gene expression. In details, three proteins Rep1, Rep2, and Faf1 which belong to the partitioning system control the copy number of plasmid by repressing expression of Flp1 protein. The amount of three repressors is in proportion to the 2-micron plasmid copy number, hence, expression of Flp1 is suppressed when the plasmid copy number is high, and if the plasmid copy number is small, Flp1 expression is induced. Therefore, expression of endogenous Flp1 is not relevant to the condition that the cell has to face. However, expression of Flp1 on an extra-plasmid transformed is segregated to the number and regulation of endogenous 2-micron system. The protein is considered as to be overexpressed and can be potentially beneficial when cells are challenged by DNA damaging agents.

We already prove that the lethality of *sgs1Δmus81<sub>Δ120N</sub>* was rescued by further deletion of Rad52, a key homologous recombination mediator in homologous recombination in budding yeast, indicating that the cellular defect related to dysfunction of Mus81 lacking N-terminal region was caused by the accumulation of unprocessed toxic recombination intermediates [26]. Therefore, the suppression of cellular defect of *sgs1Δmus81<sub>Δ100N</sub>* by Flp1 overexpression should be derived from the ability to reduce the late recombination intermediates accumulated. This notion is further supported by the result that overexpression of Flp1 was unable to rescue the cellular defect caused by the absence of Rad52 which functions upstream of Mus81 complex. The possibility that Flp1 can participate in processing toxic intermediates arising when cells are faced with DNA lesions was also supported by the observation that the overexpression of Flp1 can partially rescue the HU sensitivity of *mus81Δ* mutant at non-permissive temperature (37 °C). At 37 °C, a condition that the cells grow and divide rapidly, HU treatment induces quickly high accumulation of toxic intermediates that only Sgs1 activity could not cope with efficiently, representing by the highly drug-sensitive level of the cells (Figure 5). This excessive accumulation of recombination intermediates could lead to a condition that the cells may activate Flp1 function in mediating the removal of the toxic intermediates besides Sgs1, showing the partially resistant capability to drug (Figure 5).

Until now, the reason of 2-micron plasmid presence inside yeast cell has not been clearly defined, hence, identifying Flp1 as the suppressor of Mus81 partial dysfunction raises the possible explanation of advantage of readily containing this plasmid inside the cells as a backup system. As discussed above, we believe that Flp1 overexpression suppressed the cellular defects related to the loss of Mus81 function caused by the loss of N-terminal region via procession of toxic recombination intermediates. Although Flp1 tetramers might bind DNA in a configuration similar from recombination repair intermediates, Flp1 is a site-specific recombinase requiring a specific DNA sequence to induce DNA cleavage and recombination. It has not been completely clear that the

Ffp1 overexpression suppressor effect depends on its enzymatic activities, pointing to an involvement of DNA cleavage and recombination, or its potential function is perhaps merely structural. Therefore, Ffp1 should be further investigated for its possible function in resolving homologous recombination intermediates generated when cells try to repair DNA damages.

### Conflict of interest

All authors declare no conflicts of interest in this paper.

### References

1. Kaliraman V, Mullen JR, Fricke WM, et al. (2001) Functional overlap between Sgs1-Top3 and the Mms4-Mus81 endonuclease. *Genes Dev* 15: 2730–2740.
2. Nishino T, Komori K, Ishino Y, et al. (2003) X-ray and biochemical anatomy of an archaeal XPF/Rad1/Mus81 family nuclease: similarity between its endonuclease domain and restriction enzymes. *Structure* 11: 445–457.
3. Blais V, Gao H, Elwell CA, et al. (2004) RNA interference inhibition of Mus81 reduces mitotic recombination in human cells. *Mol Biol Cell* 15: 552–562.
4. Newman M, Murray-Rust J, Lally J, et al. (2005) Structure of an XPF endonuclease with and without DNA suggests a model for substrate recognition. *EMBO J* 24: 895–905.
5. Boddy MN, Lopez-Girona A, Shanahan P, et al. (2000) Damage tolerance protein Mus81 associates with the FHA1 domain of checkpoint kinase Cds1. *Mol Cell Biol* 20: 8758–8766.
6. Interthal H, Heyer WD (2000) MUS81 encodes a novel helix-hairpin-helix protein involved in the response to UV- and methylation-induced DNA damage in *Saccharomyces cerevisiae*. *Mol Gen Genet* 263: 812–827.
7. Abraham J, Lemmers B, Hande MP, et al. (2003) Eme1 is involved in DNA damage processing and maintenance of genomic stability in mammalian cells. *EMBO J* 22: 6137–6147.
8. Bastin-Shanower SA, Fricke WM, Mullen JR, et al. (2003) The mechanism of Mus81-Mms4 cleavage site selection distinguishes it from the homologous endonuclease Rad1-Rad10. *Mol Cell Biol* 23: 3487–3496.
9. Chen XB, Melchionna R, Denis CM, et al. (2001) Human Mus81-associated endonuclease cleaves Holliday junctions *in vitro*. *Mol Cell* 8: 1117–1127.
10. Boddy MN, Gaillard PH, McDonald WH, et al. (2001) Mus81-Eme1 are essential components of a Holliday junction resolvase. *Cell* 107: 537–548.
11. Gaillard PH, Noguchi E, Shanahan P, et al. (2003) The endogenous Mus81-Eme1 complex resolves Holliday junctions by a nick and counternick mechanism. *Mol Cell* 12: 747–759.
12. Constantinou A, Chen XB, McGowan CH, et al. (2002) Holliday junction resolution in human cells: two junction endonucleases with distinct substrate specificities. *EMBO J* 21: 5577–5585.
13. Doe CL, Ahn JS, Dixon J, et al. (2002) Mus81-Eme1 and Rqh1 involvement in processing stalled and collapsed replication forks. *J Biol Chem* 277: 32753–32759.
14. Whitby MC, Osman F, Dixon J (2003) Cleavage of model replication forks by fission yeast Mus81-Eme1 and budding yeast Mus81-Mms4. *J Biol Chem* 278: 6928–6935.
15. Gaskell LJ, Osman F, Gilbert RJ, et al. (2007) Mus81 cleavage of Holliday junctions: a failsafe for processing meiotic recombination intermediates? *EMBO J* 26: 1891–1901.

16. Schwartz EK, Wright WD, Ehmsen KT, et al. (2012) Mus81-Mms4 functions as a single heterodimer to cleave nicked intermediates in recombinational DNA repair. *Mol Cell Biol* 32: 3065–3080.
17. Matos J, Blanco MG, Maslen S, et al. (2011) Regulatory control of the resolution of DNA recombination intermediates during meiosis and mitosis. *Cell* 147: 158–172.
18. Gallo-Fernandez M, Saugar I, Ortiz-Bazan MA, et al. (2012) Cell cycle-dependent regulation of the nuclease activity of Mus81-Eme1/Mms4. *Nucleic Acids Res* 40: 8325–8335.
19. Princz LN, Wild P, Bittmann J, et al. (2017) Dbf4-dependent kinase and the Rtt107 scaffold promote Mus81-Mms4 resolvase activation during mitosis. *EMBO J* 36: 664–678.
20. Saugar I, Jimenez-Martin A, Tercero JA (2017) Subnuclear Relocalization of Structure-Specific Endonucleases in Response to DNA Damage. *Cell Rep* 20: 1553–1562.
21. Mullen JR, Kaliraman V, Ibrahim SS, et al. (2001) Requirement for three novel protein complexes in the absence of the Sgs1 DNA helicase in *Saccharomyces cerevisiae*. *Genetics* 157: 103–118.
22. Chu WK, Hickson ID (2009) RecQ helicases: multifunctional genome caretakers. *Nat Rev Cancer* 9: 644–654.
23. Fabre F, Chan A, Heyer WD, et al. (2002) Alternate pathways involving Sgs1/Top3, Mus81/Mms4, and Srs2 prevent formation of toxic recombination intermediates from single-stranded gaps created by DNA replication. *Proc Natl Acad Sci U S A* 99: 16887–16892.
24. Ciccia A, Constantinou A, West SC (2003) Identification and characterization of the human mus81-eme1 endonuclease. *J Biol Chem* 278: 25172–25178.
25. Kang MJ, Lee CH, Kang YH, et al. (2010) Genetic and functional interactions between Mus81-Mms4 and Rad27. *Nucleic Acids Res* 38: 7611–7625.
26. Thu HPT, Nguyen TA, Munashingha PR, et al. (2015) A physiological significance of the functional interaction between Mus81 and Rad27 in homologous recombination repair. *Nucleic Acids Res* 43: 1684–1699.
27. Chavdarova M, Marini V, Sisakova A, et al. (2015) Srs2 promotes Mus81-Mms4-mediated resolution of recombination intermediates. *Nucleic Acids Res* 43: 3626–3642.
28. Sisakova A, Altmannova V, Sebesta M, et al. (2017) Role of PCNA and RFC in promoting Mus81-complex activity. *BMC Biol* 15: 90.
29. Sebesta M, Urulangodi M, Stefanovic B, et al. (2017) Esc2 promotes Mus81 complex-activity via its SUMO-like and DNA binding domains. *Nucleic Acids Res* 45: 215–230.
30. Keyamura K, Arai K, Hishida T (2016) Srs2 and Mus81-Mms4 Prevent Accumulation of Toxic Inter-Homolog Recombination Intermediates. *PLoS Genet* 12: e1006136.
31. Ghamrasni SE, Cardoso R, Li L, et al. (2016) Rad54 and Mus81 cooperation promotes DNA damage repair and restrains chromosome missegregation. *Oncogene* 35: 4836–4845.
32. Ii M, Ii T, Brill SJ (2007) Mus81 functions in the quality control of replication forks at the rDNA and is involved in the maintenance of rDNA repeat number in *Saccharomyces cerevisiae*. *Mutat Res* 625: 1–19.
33. Fitcher AB (1988) The 2 micron circle plasmid of *Saccharomyces cerevisiae*. *Yeast* 4: 27–40.
34. Chan KM, Liu YT, Ma CH, et al. (2013) The 2 micron plasmid of *Saccharomyces cerevisiae*: a miniaturized selfish genome with optimized functional competence. *Plasmid* 70: 2–17.
35. Ahn YT, Wu XL, Biswal S, et al. (1997) The 2microm-plasmid-encoded Rep1 and Rep2 proteins interact with each other and colocalize to the *Saccharomyces cerevisiae* nucleus. *J Bacteriol* 179: 7497–7506.

36. Velmurugan S, Ahn YT, Yang XM, et al. (1998) The 2 micrometer plasmid stability system: analyses of the interactions among plasmid- and host-encoded components. *Mol Cell Biol* 18: 7466–7477.
37. Velmurugan S, Yang XM, Chan CS, et al. (2000) Partitioning of the 2-microm circle plasmid of *Saccharomyces cerevisiae*. Functional coordination with chromosome segregation and plasmid-encoded rep protein distribution. *J Cell Biol* 149: 553–566.
38. Ghosh SK, Hajra S, Paek A, et al. (2006) Mechanisms for chromosome and plasmid segregation. *Annu Rev Biochem* 75: 211–241.
39. Andrews BJ, Proteau GA, Beatty LG, et al. (1985) The FLP recombinase of the 2 micron circle DNA of yeast: interaction with its target sequences. *Cell* 40: 795–803.
40. Gronostajski RM, Sadowski PD (1985) The FLP recombinase of the *Saccharomyces cerevisiae* 2 microns plasmid attaches covalently to DNA via a phosphotyrosyl linkage. *Mol Cell Biol* 5: 3274–3279.



AIMS Press

© 2018 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)