

# Multiparametric magnetic resonance imaging of prostate cancer

Dear Sir,

We read the interesting article by Hedgire *et al.*, titled "Multiparametric magnetic resonance imaging of prostate cancer,"<sup>[1]</sup> which was published in the August 2012 issue of the journal. The study included fairly important information which was very useful for us. However, we would like to make a few contributions.

Mucinous-type lesions are rare adenocarcinomas of the prostate gland and tend to show hyperintensity on T2-weighted images. Thus, mucinous tumors may be missed in T2-weighted images. Likewise, mucinous lesions may be missed in diffusion-weighted imaging (DWI) since high apparent diffusion coefficient (ADC) values may be obtained for mucinous tumors.<sup>[2]</sup> We believe that MR spectroscopy may provide more realistic results in determining these lesions.

In our experience of prostate MRI and histopathologic results, we did not observe any prostate cancer focus at the hematoma site in the peripheral zone on prostate MR examination obtained 2-3 weeks after prostate biopsies. However, we detected cancer focus at the non-hemorrhagic sites in the peripheral zone. In our opinion, the reason for this is that hematoma cannot dissect the tumor composed of tightly packed cells. Therefore, we recommend that non-hemorrhagic sites in the peripheral zone on

T1-weighted images should especially be evaluated for the presence of tumor focus on the other imaging sequences.

**Ali Kemal Sivrioglu, Mehmet AK<sup>1</sup>, Kemal Kara<sup>2</sup>,  
Güner Sönmez<sup>2</sup>**

Department of Radiology, AKSAZ Military Hospital, Mugla,  
<sup>1</sup>Department of Radiology, Erzurum Maresal Cakmak Hospital,  
Erzurum, <sup>2</sup>Department of Radiology, GATA Haydarpasa  
Training Hospital, İstanbul, Turkey  
E-mail: draksivrioglu@gmail.com

## References

1. Hedgire SS, Oei TN, Mcdermott S, Cao K, Patel MZ, Harisinghani MG. Multiparametric magnetic resonance imaging of prostate cancer. *Indian J Radiol Imaging* 2012;22:160-9.
2. Sonmez G, Cuce F, Mutlu H, Incedayi M, Ozturk E, Sildiroglu O, *et al.* Value of diffusion-weighted MRI in the differentiation of benign and malign breast lesions. *Wien Klin Wochenschr* 2011;123:655-61.

### Access this article online

#### Quick Response Code:



**Website:**  
www.ijri.org

**DOI:**  
10.4103/0971-3026.116580