

Molecular Epidemiology of Methicillin-resistant *Staphylococcus aureus* in Children: Comparison between Cystic Fibrosis and Non-cystic Fibrosis Isolates

Sir,

We have previously reported, elsewhere and separately, the molecular and phenotypical characteristics of methicillin-resistant *Staphylococcus aureus* (MRSA) isolates from cystic fibrosis patients and from non-cystic fibrosis patients.^[1,2] We aimed to perform a molecular comparison between the isolates from the two groups, using pulsed field gel electrophoresis (PFGE).

Thirty seven MRSA isolates were included, nine from cystic fibrosis patients (respiratory secretion), and 30 from non-cystic fibrosis, hospitalized patients (from

various anatomical sites). One patient with cystic fibrosis and three patients without cystic fibrosis had two isolates included each. Of the 39 isolates, one carried SCCmec type I, three SCCmec type II, 19 SCCmec type III, and 16 SCCmec type IV. DNA restriction patterns after digestion were analysed by PFGE, which was performed as described previously.^[3] Results were compared using the GelComparII software (Applied Maths, Kortrijk, Belgium), based on published criteria.^[4]

Cystic fibrosis isolates were similar to non-cystic fibrosis isolates. We observed clustering within SCCmec types (notably types III and IV, that were more numerous), but not within groups of isolates (cystic fibrosis versus non-cystic fibrosis). Most type III isolates had patterns typical of the Brazilian clone, whereas the majority of type IV isolates were at least similar to the SCCmec IV pediatric clone. These findings suggest that the local of acquisition of MRSA isolates could be the same (probably the hospital) for most of the cystic fibrosis and non-cystic fibrosis patients. Prospective studies with a higher number of isolates are necessary to clarify the details of the epidemiological relationship between MRSA isolates in these patients.

**Marcelo Jenne Mimica^{1,2},
Eitan Naaman Berezin¹, Neiva Damaceno¹,
Rozane Bigelli Carvalho²**

Departments of ¹Pediatrics and ²Pathology,
Santa Casa School of Medicine, Rua Cesário Mota Jr.,
61, São Paulo, Brazil

Address for correspondence: Dr. Marcelo Mimica,
E-mail: mjmimica@hotmail.com

REFERENCES

1. Mimica MJ, Berezin EN, Carvalho RB. Healthcare associated PVL-negative methicillin-resistant *Staphylococcus aureus* with SCCmec type IV. *Pediatr Infect Dis J* 2009;28:934.
2. Mimica MJ, Berezin EN, Damaceno N, Carvalho RB. SCCmec Type IV, PVL-Negative, Methicillin-Resistant *Staphylococcus aureus* in Cystic Fibrosis Patients from Brazil. *Curr Microbiol* 2011;62:388-90.
3. Hultén KG, Kaplan SL, Gonzalez BE, Hammerman WA, Lamberth LB, Versalovic J, *et al.* Three-years surveillance of community onset health care-associated *Staphylococcus aureus* infections in children. *Pediatr Infect Dis J* 2006;25:349-53.
4. Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, *et al.* Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;33:2233-9.

Access this article online

Quick Response Code:



Website:
www.jlponline.org

DOI:
10.4103/0974-2727.98683