## Images in Gastroenterology

## A young female with recurrent biliary pain

Rajesh Puri, Narendra S. Choudhary, Randhir Sud

Institute of Digestive and Hepatobiliary Sciences, Medanta, The Medicity, Gurgaon, Haryana, India

A 27-year-old female from Rajasthan presented with history of recurrent biliary colic for past 6 months. She was investigated at another hospital. Her liver function test was normal. Ultrasound of the abdomen was done to exclude gall stones; it showed normal gall bladder and common bile duct. Magnetic resonance cholangiopancreatography (MRCP) showed normal common bile duct diameter and there were no filling defects. An endoscopic ultrasound (EUS) was done to look for microlithiasis. EUS revealed a linear structure in gallbladder with a central hypoechoic defect as shown in Figure 1. The structure exhibited movements during examination consistent with the presence of adult roundworm, *Ascaris lumbricoides* within the gallbladder. The patient was given albendazole and cholecystectomy was done. It confirmed presence of ascaris worm inside the gallbladder. She is asymptomatic at 1 year of follow-up.

Ascaris lumbricoides is the most common helminthic infection of the world and approximately 25% of population is affected. It is transmitted via the fecal-oral route, mainly from ingestion of water or food (vegetables or fruits) contaminated with A. lumbricoides eggs. Infection remains asymptomatic in majority of the infected persons. Symptoms are infrequent and may be caused by larvae migration in circulation or by adult worms migration into pancreatobiliary system.[1] Adult worm may cause biliary colic, cholecystitis, cholangitis, pancreatitis, hepatic abscess, intestinal obstruction and acute appendicitis.<sup>[2-4]</sup> Diagnosis of biliary ascariasis in nonendemic area requires high index of suspicion. In ultrasound or endoscopic ultrasound, Ascaris lumbricoides appears as long echogenic structure without producing shadow effect, as it may appear like a coil or as echogenic structure with central anechoic linear defect. In gall bladder, its zig-zag movement may be seen. [5,6] Gall bladder ascariasis is rare, in a series of 500 hepatobiliary pancreatic diseases due to Ascaris; only 8 (1.6%) patients had worm in gall

Access this article online	
	Quick Response Code
Website:	
www.jdeonline.in	<u> </u>
	(A)
	48222
DOI:	
10.4103/0976-5042.132441	回逐步被




Figure 1: EUS image showing long linear echogenic structure with linear central hypoechoic defect and coiling on itself in gall bladder

bladder.<sup>[2]</sup> Gall bladder worm generally need cholecystectomy, although conservative management (antihelminthic therapy) alone may work for some patients.<sup>[6,7]</sup>

## References

- Elliot DE. Intestinal infections by parasitic worms. In: Feldman M, Friedman LS, Brandt LJ, editors. Sleisenger and Fordtran's Gastrointestinal and Liver Diseases. 9th ed. p. 1921-39.
- Khuroo MS, Zargar SA, Mahajan R. Hepatobiliary and pancreatic ascariasis in India. Lancet 1990;335:1503-6.
- Coskun A, Ozcan N, Durak AC, Tolu I, Güleç M, Turan C. Intestinal ascariasis as a cause of bowel obstruction in two patients: Sonographic diagnosis. J Clin Ultrasound 1996;24:326-8.
- Misra SP, Dwivedi M, Misra V, Singh AP, Agarwal VK. Preoperative sonographic diagnosis of acute appendicitis caused by Ascaris lumbricoides. J Clin Ultrasound 1999;27:96-7.
- Khuroo MS, Zargar SA, Yattoo GN, Dar MY, Javid G, Khan BA, et al. Sonographic findings in gallbladder ascariasis. J Clin Ultrasound 1992;20:587-91.
- Javid G, Wani N, Gulzar GM, Javid O, Khan B, Shah A. Gallbladder ascariasis: Presentation and management. Br J Surg 1999;86:1526-7.
- Cha DY, Song IK, Choi HW, Chung EA, Shin BS, Song YW, et al. Successful elimination of Ascaris lumbricoides from the gallbladder by conservative medical therapy. J Gastroenterol 2002;37:758-60.

**How to cite this article:** Puri R, Choudhary NS, Sud R. A young female with recurrent biliary pain. J Dig Endosc 2013;4:138.

Source of Support: Nil, Conflict of Interest: None declared.

## Address for correspondence

Dr. Rajesh Puri, Therapeutic Endoscopist and Endosonologist, Institute of Digestive and Hepatobiliary Sciences, Medanta, The Medicity, Gurgaon - 122 001, Haryana, India. E-mail: purirajesh69@gmail.com