

The influence of type 2 diabetes mellitus on salivary matrix metalloproteinase-8 levels and periodontal parameters: A study in an Indian population

Namita Gupta¹, Narinder Dev Gupta¹, Akash Gupta², Lata Goyal¹, Sagar Garg³

¹Department of Periodontics and Community Dentistry, Dr. Z. A. Dental College, Aligarh Muslim University, Aligarh, Uttar Pradesh, India,

²Department of Biochemistry, Subharti Medical College, Meerut, Uttar Pradesh, India,

³Department of Periodontics, Tamil Nadu Government Medical College, Chennai, Tamil Nadu, India

Correspondence: Dr. Akash Gupta
Email: akash_inspace@yahoo.com

ABSTRACT

Objective: Although many studies reported more severe periodontal disease and the existing proinflammatory conditions in patients with diabetes but only few have examined the effect of type 2 diabetes mellitus (DM) on salivary matrix metalloproteinase-8 (MMP-8) level and other periodontal parameters. This study aims to evaluate the effect of type 2 DM on salivary MMP-8 levels and periodontal parameters, which might be useful in monitoring periodontal disease in diabetes. **Materials and Methods:** A total of 90 subjects were selected for the study and were divided into three groups: Group I included 30 healthy subjects; Group II included 30 subjects without type 2 DM but with chronic periodontitis, and Group III included 30 subjects with type 2 DM and chronic periodontitis. Periodontal parameters such as plaque index (PI), gingival index (GI), pocket probing depth (PPD), and clinical attachment level (CAL) were taken. The salivary MMP-8 level was estimated by Quantikine Human total MMP-8 immunoassay kit using ELISA method. **Results:** The mean value of the salivary MMP-8 of Group III was highest followed by Group II and Group I, the least. The other periodontal parameters PI, GI, PPD, CAL, was comparatively highest for Group III. **Conclusion:** This study suggests that diabetes is associated with an increased prevalence, extent, and severity of periodontitis. Furthermore, the increased levels of MMP-8 indicate the influence of diabetes on their salivary concentration.

Key words: Chronic periodontitis, matrix metalloproteinase-8, periodontal parameters, saliva, type 2 diabetes mellitus

INTRODUCTION

Mankind has been tormented by diseases of oral cavity since time immemorial. Periodontitis is a multifactorial inflammatory disease affecting the periodontal tissues, which is characterized by progressive destruction of the tissues supporting the tooth, and finally the outcome of untreated periodontitis is tooth loss.^[1] The role of host defense response is usually most important aspect in periodontitis mainly because it can determine both disease progression along with severity.^[2] Diabetes mellitus (DM) is one such wide existing endocrine disorder, which is known to alter this delicate balance thus imposing exaggerated tissue damage whenever challenged with a disease like periodontitis.^[3,4] Periodontal disease has been proposed as the sixth

complication of diabetes.^[5] Diabetes itself does not trigger periodontitis nevertheless it predisposes and speeds up the irreversible periodontal destruction. Individuals with diabetes have increased defective polymorphonuclear function, increased collagenase

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Gupta N, Gupta ND, Gupta A, Goyal L, Garg S. The influence of type 2 diabetes mellitus on salivary matrix metalloproteinase-8 levels and periodontal parameters: A study in an Indian population. *Eur J Dent* 2015;9:319-23.

DOI: 10.4103/1305-7456.163222

production, altered collagen and glycosaminoglycan synthesis, deregulation in cytokine production, and the formation of advanced glycation end products (AGEs);^[6] and they contribute to tissue destruction by inducing the release of cytokines through a multistep process, which ultimately leads to formation of more of matrix metalloproteinases (MMPs).^[7]

Enzymes play a crucial role in the regulation of connective tissue destruction in periodontitis, most notably MMPs. Under normal physiological conditions, the expression of MMPs is considerably low, and these low levels are associated with healthy connective tissue remodeling.^[8] In pathologic conditions such as periodontitis,^[9] oral cysts,^[10] oral cancer,^[11] and atherosclerosis;^[12] the level of MMP expression enhances significantly, resulting in aberrant connective tissue destruction, hence leads to the elevation of MMP levels in saliva. MMPs are proteolytic enzymes belonging to zinc protease superfamily and are involved in physiological degradation of extracellular matrix protein and basement membrane. There are several groups of MMPs: Collagenase (MMP 1, 8, 13), gelatinase (MMP 2, 9), stromelysin (MMP 3, 10, 11), matrilysins (MMP 7), and other membrane associated MMPs (2, 3). In the present era, of periodontal medicine and therapeutics, traditional diagnostic aids such as probing depth and attachment loss have certain limitations. To overcome these challenges, researchers are trying to find out new biomarkers that can enable the clinicians in assessing and diagnosing the patient's condition better.^[13] Research shows MMP-8, a neutrophil collagenase, as a potential biomarker in chronic periodontitis, and it is highly correlated with bleeding on probing, probing pocket depth, and attachment loss.^[14,15] It has the unique ability to decompose Types I and III collagen.^[16] This study was aimed at understanding the contribution of type 2 DM in the pathogenesis of periodontitis. In order to achieve this goal, the periodontal parameters, and the levels of salivary MMP-8 were measured.

MATERIALS AND METHODS

Patient selection

The present study was conducted in the Department of Periodontology of Dr. Z. A Dental College, Aligarh Muslim University, Aligarh, India. This study was approved by the Institutional Review Board (Department of Periodontics and Community Dentistry, Aligarh Muslim University, Aligarh). All patients were individually informed about the study, and informed consent forms were signed in full accordance with the Helsinki Declaration.

Groups

A total of 90 (40 males and 41 females) individuals in the age range of 45–55 years and having at least 20 teeth were taken into the study and were further divided into three groups: Group I: Healthy subjects, Group II: Chronic periodontitis patients without type 2 DM, and Group III: Chronic periodontitis with type 2 DM. Each group consisted of 30 subjects.

Inclusion criteria

Group I

Periodontally and systemically healthy individuals with no apparent signs of clinical inflammation.

Group II

Subjects with no systemic disease but clinically diagnosed with moderate to severe chronic periodontitis (e.g., at least two or more interproximal sites with clinical attachment level (CAL) of ≥ 4 mm, or two or more interproximal sites with pocket probing depth (PPD) of ≥ 5 mm, not on the same tooth).^[17]

Group III

Subjects with type 2 DM with current metabolic control above the normal range (glycated hemoglobin [HbA1c] test $> 6\%$) and clinically diagnosed with moderate to severe chronic periodontitis (e.g., at least two or more interproximal sites with CAL of ≥ 4 mm, or two or more interproximal sites with PPD of ≥ 5 mm, not on the same tooth).

Exclusion criteria

Exclusion criteria included smokers, pregnant and lactating women, and patients with acute or chronic medical disorders. Patients who had undergone any surgery in past 6 months, subjects undergoing orthodontic treatment were not included in the study.

Clinical measurements

Periodontal disease status of all patients was evaluated by measurement of plaque index (PI),^[18] gingival index (GI),^[19] PPD, and CAL. To rule out any bias, same instruments (University of North Carolina-15 probe [UNC-15], Hu-Freidy's USA) were used by the same examiner. Intraexaminer reliability of index reproducibility was assessed twice during the study in 10 patients by repeating the measurements (k-score for each measurement was never lower than 0.76). The pocket depth is the distance between the base of the pocket and the gingival margin. The distance was calculated with the UNC-15 probe held parallel to the vertical axis of the tooth. The pocket depth was measured at four sites per tooth. The CAL was calculated - It is the distance between the bases of the

pocket to cemento-enamel junction. All patients were checked for their diabetic status by testing for HbA1c levels (by D-10 hemoglobin analyzer from Bio Rad). The same laboratory is used for testing for all the patients.

Determination of matrix metalloproteinase-8 levels in saliva

Saliva sample collection

The clinical parameters described above were recorded, and saliva samples necessary for estimation of MMP-8 were taken from all groups. Subjects were told not to eat or drink 2 h prior to the examination. They were seated comfortably, then asked to rinse and 3 ml of unstimulated whole expectorated saliva was collected from each subject and placed into sterile 5 ml saliva collecting tubes according to the method described by Navazesh.^[20] The collected saliva was centrifuged immediately to remove cell debris (1000 ×g for 10 min at 4°C). Then the supernatant was removed, and the sample was stored in small aliquots at -80°C until analysis.

MMP-8 levels Estimation

Salivary MMP-8 levels were estimated for each subject using Quantikine Human total MMP-8 immunoassay kit using ELISA method, provided by R&D Systems, (Minneapolis USA). Absorbance was read at 450 nm, with the correction wavelength set at 540 or 570 nm. The minimum detectable dose of MMP-8 ranged from 0.01 to 0.06 ng/ml.

Statistical analysis

Continuous data were summarized as the mean ± standard deviation while discrete (categorical) in percentage. Continuous variables were compared by one-way analysis of variance (ANOVA), and the significance of the mean difference between the groups was done by Tukey's *post-hoc* test. Categorical variables were compared by Chi-square test. The statistical significance of correlations among MMP-8 and periodontal parameters was determined using spearman rank correlation coefficients. A two-sided ($\alpha = 2$) $P < 0.05$ was considered as statistically significant. All analyses were performed on STATISTICA (Windows version 6.0 for 2001. StatSoft, Inc.,)

RESULTS

The mean values for the clinical parameters and demographic data of all the three groups are given in Table 1. No significant difference between the mean age was found among the three groups ($P > 0.05$). Similarly, no significant difference was found in the proportion of genders (M/F) between the three groups ($P > 0.05$).

A comparison among all three groups showed a significant difference in periodontal parameters ($P < 0.001$). The mean PI, GI, PPD, CAL values of Groups II and III were significantly higher than those of Group I ($P < 0.001$), with the highest values in Group III.

MMP-8 levels: ANOVA revealed the mean values of salivary MMP-8 of Group II (381.60 ± 57.66) and III (458.36 ± 45.52) were significantly higher ($P < 0.001$) than Group I (205.04 ± 89.14), with the highest being in Group III. Further, Tukey test revealed that the mean MMP-8 levels of Groups I and III were significantly ($P < 0.001$) different as compared to Group I [Figure 1]. The data of Table 2 revealed a highly significant positive correlation (r) between MMP-8 and periodontal parameters in the entire sample.

DISCUSSION

Current studies are in a support that in patients with diabetes, the incidence and severity of periodontitis are much greater as compared to nondiabetics,^[21] to the extent that periodontitis has been called the "sixth complication of diabetes."^[5] The reason being a combination of various factors that render the host resistance weakened against periodontal pathogens. There is decreased PMN chemotaxis, an increase in the level of enzymes such as collagenase, elastase, and the decrease in collagen formation by fibroblasts that are responsible for the periodontal destruction.^[22]

Table 1: Demographic data and clinical parameters (mean ± SD) of study groups

Parameters	Group I (n=30)	Group II (n=30)	Group III (n=30)	P
Age (years)	44.87±4.34	46.80±4.92	48.96±3.75	0.427*
Gender (%)				
Male	16 (53.3)	16 (53.3)	17 (56.7)	0.956*
Female	14 (46.7)	14 (46.7)	13 (43.3)	
PI	0.54±0.10	1.57±0.39	2.30±0.23	<0.001***
GI	0.52±0.10	2.06±0.18	2.45±0.17	<0.001***
PPD	2.10±0.24	5.34±0.24	6.06±0.37	<0.001***
CAL	0.60±0.07	5.30±0.20	5.97±0.42	<0.001***
HbA1c	4.78±0.42	4.90±0.69	7.03±0.65	-

Data are expressed as mean±SD. Group I (control): Periodontally and systemically healthy individuals with no apparent signs of clinical inflammation (n=30), Group II (chronic periodontitis without type 2 DM): Subjects with no systemic disease but clinically diagnosed with moderate to severe chronic periodontitis (e.g., at least two or more interproximal sites with CAL of ≥4 mm, or two or more interproximal sites with PPD of ≥5 mm, not on the same tooth) (n=30), Group III (chronic periodontitis with type 2 DM): Subjects with type 2 DM with current metabolic control above the normal range (HbA1c test >6%) and clinically diagnosed with moderate to severe chronic periodontitis (e.g., at least two or more interproximal sites with CAL of ≥4 mm, or two or more interproximal sites with PPD of ≥5 mm, not on the same tooth) (n=30). *Not significant, ***Highly significant. PI: Plaque index, GI: Gingival index, PPD: Probing pocket depth, CAL: Clinical attachment level, HbA1c: Glycated hemoglobin, DM: Diabetes mellitus




Figure 1: Salivary MMP-8 levels (mean \pm SD) of three groups. Group I (control), Group II (chronic periodontitis without type 2 DM, Group III (chronic periodontitis with type 2 DM). In Group III, the levels of MMP-8 were significantly increased as compared to group II and group I. a) Statistically significant difference as compared between group I and II ($P < 0.001$). b) Statistically significant difference as compared between groups II and III ($P < 0.001$)

Table 2: Correlation between salivary MMP-8 and periodontal parameters

Parameters	Correlation (r) value	P^*
PI	0.74	0.000
GI	0.72	0.000
PPD	0.80	0.000
CAL	0.78	0.000

*Correlation is significant at the 0.01 level (two-tailed). PI: Plaque index, GI: Gingival index, PPD: Probing pocket depth, CAL: Clinical attachment level, MMP-8: Matrix metalloproteinase-8

Our results corroborate the findings of several studies reporting a positive association between diabetes and periodontal disease, which reported a greater risk of attachment loss and bone loss in diabetic patients.^[23,24] In diabetes, the formation of collagen and glycosaminoglycan is impaired, collagenolytic activity of gingival crevice fluid is increased, which in turn causes loss of periodontal fibers, loss of the alveolar bone, and finally results in exfoliation of the teeth.^[25] Moreover, in the present study, it was found that the value of GI was higher for diabetics as compared to nondiabetic patients, which concur with the finding reported by Cutler *et al.*^[26] and Ervasti *et al.*^[27] who reported that poorly controlled diabetics had mean GI score of 77.9 ± 19.2 as compared to 61.9 ± 25.1 in the control group, and it was statistically significant. This may be due to the metabolic imbalances in the tissue which can lower the resistance to infection in diabetic individuals. Decrease in response to inflammation may also be due to impaired neutrophil chemotaxis in diabetic patients.^[28]

MMP-8 has been considered as a key biomarker in chronic periodontitis.^[29] It is a major collagenolytic

metalloproteinase in gingival tissues and oral fluids, and its elevated level is associated with the severity of periodontal inflammation.^[14] In this study, the quantitative analysis of MMP-8 showed that MMP-8 levels were significantly increased in patients with type 2 diabetic (Group III) as compared to nondiabetics (Group II), although both having generalized chronic periodontitis. Thus, the difference between MMP-8 levels in Groups II and III was because of periodontal destruction, which is more in type 2 diabetes. A possible justification pertaining to this could be that in diabetes, collagen crosslinking occurs due to the formation of AGE that could induce a cycle of cytokine up-regulation after binding to macrophage receptors. This results in an elevated expression of MMP in diabetic periodontal tissues.^[30] Comparing the rise of MMP-8 levels, which shows a higher mean salivary MMP-8 in diabetic patients as compared to nondiabetic and control group is highly in conformity with the result obtained by Costa *et al.*^[31] who observed highest concentrations of MMP-8 in the saliva of patients with diabetes. Our result also comes in agreement with the results of Collin *et al.*,^[32] Kumar *et al.*,^[33] which found a rise in MMP-8 levels as compared to nondiabetic controls in saliva. Another study by Rathnayake *et al.*^[34] also revealed that patients with diabetes had a 3 times higher ratio of the MMP-8/TIMP-1 and twice as high concentration of MMP-8. Furthermore, in the present study, the MMP-8 concentration had a significant positive correlation with periodontal indices. This indicates that periodontal disease severity was increased with the values of MMP-8, and it has a strong association between periodontal parameters and MMP-8.

Unstimulated saliva was used as the sample for study. Using saliva for sampling has certain advantages such as the easy collection, noninvasive, and rapid procedure; moreover, it can be collected from all the sites in the mouth giving an overall assessment of disease status and severity.^[35] Furthermore, its measurement does not require any special equipment and expertise.

Diabetes is certainly one of the biggest and most serious risk factor in development and progression of periodontitis as the latter is significantly aggravated in patients suffering from diabetes having long-term hyperglycemia.^[36] Although a lot has been achieved in establishing the association between the accelerated periodontal disease and diabetes, but more breakthroughs are needed. While the finding of the study could be applied in most instances, there were some limitations of this study, such as the sample size was small. Thus, studies with larger sample size should be done so that various associated risk factors could be studied in detail.

CONCLUSION

In summary, the current study suggests that the diabetes is a major risk factor that exerts a negative effect on periodontal tissues, which contribute to an enhanced susceptibility to periodontitis. Furthermore, the increased levels of MMP-8 in saliva clearly point toward active periodontitis in diabetic patients. This opens up new possibilities in future, specific immunoassays targeting MMPs, especially MMP-8 or collagenase-2 from oral fluids, could prove out to be useful in monitoring the course of periodontitis in patients with diabetes.

Acknowledgments

The authors are grateful to all the participants in this study and to the Department of Periodontics and Community Dentistry, Dr. Z. A. Dental College, Aligarh Muslim University, Aligarh, India. All the authors have significantly contributed in the study.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Nunn ME. Understanding the etiology of periodontitis: An overview of periodontal risk factors. *Periodontol* 2000;32:11-23.
- Seymour GJ, Gemmell E, Reinhardt RA, Eastcott J, Taubman MA. Immunopathogenesis of chronic inflammatory periodontal disease: Cellular and molecular mechanisms. *J Periodontol Res* 1993;28:478-86.
- Shlossman M, Knowler WC, Pettitt DJ, Genco RJ. Type 2 diabetes mellitus and periodontal disease. *J Am Dent Assoc* 1990;121:532-6.
- Mealey BL, Ocampo GL. Diabetes mellitus and periodontal disease. *Periodontol* 2000;44:127-53.
- Löe H. Periodontal disease. The sixth complication of diabetes mellitus. *Diabetes Care* 1993;16:329-34.
- Schmidt AM, Hori O, Cao R, Yan SD, Brett J, Wautier JL, *et al.* RAGE: A novel cellular receptor for advanced glycation end products. *Diabetes* 1996;45:577-80.
- Ryan ME, Ramamurthy NS, Sorsa T, Golub LM. MMP-mediated events in diabetes. *Ann N Y Acad Sci* 1999;878:311-34.
- Nagase H, Woessner JF Jr. Matrix metalloproteinases. *J Biol Chem* 1999;274:21491-4.
- Bertoldi C, Lalla M, Pradelli JM, Cortellini P, Lucchi A, Zaffe D. Risk factors and socioeconomic condition effects on periodontal and dental health: A pilot study among adults over 50 years of age. *Eur J Dent* 2013;7:336-46.
- Wahlgren J, Maisi P, Sorsa T, Sutinen M, Tervahartiala T, Pirilä E, *et al.* Expression and induction of collagenases (MMP-8 and -13) in plasma cells associated with bone-destructive lesions. *J Pathol* 2001;194:217-24.
- Moilanen M, Pirilä E, Grénman R, Sorsa T, Salo T. Expression and regulation of collagenase-2 (MMP-8) in head and neck squamous cell carcinomas. *J Pathol* 2002;197:72-81.
- Krupinski J, Turu MM, Font MA, Ahmed N, Sullivan M, Rubio F, *et al.* Increased tissue factor, MMP-8, and D-dimer expression in diabetic patients with unstable advanced carotid atherosclerosis. *Vasc Health Risk Manag* 2007;3:405-12.
- Fine DH, Mandel ID. Indicators of periodontal disease activity: An evaluation. *J Clin Periodontol* 1986;13:533-46.
- Giannobile WV, Beikler T, Kinney JS, Ramseier CA, Morelli T, Wong DT. Saliva as a diagnostic tool for periodontal disease: Current state and future directions. *Periodontol* 2000;50:52-64.
- Kraft-Neumärker M, Lorenz K, Koch R, Hoffmann T, Mäntylä P, Sorsa T, *et al.* Full-mouth profile of active MMP-8 in periodontitis patients. *J Periodontol Res* 2012;47:121-8.
- Sorsa T, Tjäderhane L, Konttinen YT, Laihio A, Salo T, Lee HM, *et al.* Matrix metalloproteinases: Contribution to pathogenesis, diagnosis and treatment of periodontal inflammation. *Ann Med* 2006;38:306-21.
- Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol* 2007;78:1387-99.
- Silness J, Loe H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand* 1964;22:121-35.
- Loe H, Silness J. Periodontal disease in pregnancy. I. Prevalence and severity. *Acta Odontol Scand* 1963;21:533-51.
- Navazesh M. Methods for collecting saliva. *Ann N Y Acad Sci* 1993;694:72-7.
- Cificbasi E, Kantarci A, Badur S, Issever H, Cintan S. Impact of metronidazole and amoxicillin combination on matrix metalloproteinases-1 and tissue inhibitors of matrix metalloproteinases balance in generalized aggressive periodontitis. *Eur J Dent* 2015;9:53-9.
- Mealey BL, Oates TW. American academy of periodontology. Diabetes mellitus and periodontal diseases. *J Periodontol* 2006;77:1289-303.
- Campus G, Salem A, Uzzau S, Baldoni E, Tonolo G. Diabetes and periodontal disease: A case-control study. *J Periodontol* 2005;76:418-25.
- Collin HL, Uusitupa M, Niskanen L, Kontturi-Närhi V, Markkanen H, Koivisto AM, *et al.* Periodontal findings in elderly patients with non-insulin dependent diabetes mellitus. *J Periodontol* 1998;69:962-6.
- Kiran M, Arpak N, Unsal E, Erdogan MF. The effect of improved periodontal health on metabolic control in type 2 diabetes mellitus. *J Clin Periodontol* 2005;32:266-72.
- Cutler CW, Machen RL, Jotwani R, Iacopino AM. Heightened gingival inflammation and attachment loss in type 2 diabetics with hyperlipidemia. *J Periodontol* 1999;70:1313-21.
- Ervasti T, Knuutila M, Pohjamo L, Haukipuro K. Relation between control of diabetes and gingival bleeding. *J Periodontol* 1985;56:154-7.
- Alba-Loureiro TC, Munhoz CD, Martins JO, Cerchiaro GA, Scavone C, Curi R, *et al.* Neutrophil function and metabolism in individuals with diabetes mellitus. *Braz J Med Biol Res* 2007;40:1037-44.
- Gursoy UK, Könönen E, Huuomonen S, Tervahartiala T, Pussinen PJ, Suominen AL, *et al.* Salivary type I collagen degradation end-products and related matrix metalloproteinases in periodontitis. *J Clin Periodontol* 2013;40:18-25.
- Lalla E, Lamster IB, Stern DM, Schmidt AM. Receptor for advanced glycation end products, inflammation, and accelerated periodontal disease in diabetes: Mechanisms and insights into therapeutic modalities. *Ann Periodontol* 2001;6:113-8.
- Costa PP, Trevisan GL, Macedo GO, Palioto DB, Souza SL, Grisi MF, *et al.* Salivary interleukin-6, matrix metalloproteinase-8, and osteoprotegerin in patients with periodontitis and diabetes. *J Periodontol* 2010;81:384-91.
- Collin HL, Sorsa T, Meurman JH, Niskanen L, Salo T, Rönkä H, *et al.* Salivary matrix metalloproteinase (MMP-8) levels and gelatinase (MMP-9) activities in patients with type 2 diabetes mellitus. *J Periodontol Res* 2000;35:259-65.
- Kumar MS, Vamsi G, Sriprya R, Sehgal PK. Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus. *J Periodontol* 2006;77:1803-8.
- Rathnayake N, Akerman S, Klinge B, Lundegren N, Jansson H, Tryselius Y, *et al.* Salivary biomarkers for detection of systemic diseases. *PLoS One* 2013;8:e61356.
- Miller CS, King CP Jr, Langub MC, Kryscio RJ, Thomas MV. Salivary biomarkers of existing periodontal disease: A cross-sectional study. *J Am Dent Assoc* 2006;137:322-9.
- Deshpande K, Jain A, Sharma R, Prashar S, Jain R. Diabetes and periodontitis. *J Indian Soc Periodontol* 2010;14:207-12.

Access this article online

Quick Response Code:



Website:

www.eurjdent.com