Ibnosina J Med BS

QUIZ

EKG Quiz: "Confusion EKG!"

Yousef Darrat¹, Khalid Abozguia², Fathi Idris Ali³

¹University of Kentucky, Lexington, KY, USA.
²Lancashire Cardiac Centre, Blackpool Victoria Hospital, Blackpool, UK.
³Department of Medicine-Cardiology, University Medical Center, Lebanon, TN, USA.

E-mail: fidrees100@hotmail.com

Corresponding author: Dr. Fathi Idris Ali Published: 04 October 2016 Ibnosina J Med BS 2016;8(6):193-194 Received: 24 September 2016 Accepted: 25 September 2016

This article is available from: http://www.ijmbs.org

This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

History

A 78-year-old lady was admitted with deteriorating level of consciousness over a few days. The EKG below was performed upon admission. Please examine it and answer the questions before you proceed.



Questions

1. What is the major abnormality in this EKG? Any particular diagnosis suggested?

2. Would you worry if you see Digoxin on this patient's medication list?

Ibnosina Journal of Medicine and Biomedical Sciences (2016)

Answers

1. The rhythm is sinus at about 90 bpm. The most prominent abnormality is the diffuse abnormal repolarization changes. This is primarily due to a very short QT interval. In some leads, there is no "ST segment", i.e. the T wave immediately follows the QRS complex (best seen in leads V1-V3). The calculated QT and QTc are 306 ms and 370 ms respectively. These EKG findings are very suggestive of hypercalcemia.

2. Yes. Hypercalcemia may enhance the effect of digoxin and hence can potentially predispose to digitalis toxicity (1-3).

Discussion

This lady was dry and was found to have Ca++ of 3.9 mmol/L. Hypercalcemia is the likely cause of her deteriorating level of consciousness. The family gave a history of recurrent fractures, and she was admitted to the hospital for treatment and further work up to exclude malignancy. She was found to have metastatic breast cancer.

The pattern of absent "ST segment" is very characteristic of hypercalcemia. In the recent years, a familial short QT syndrome (SQTS) has been described, and was found to be associated with increased risk of sudden cardiac death. The typical EKG appearance of that syndrome is, however, different than the one shown in this case (1-3). The discussion of that syndrome is beyond the scope of this clinical vignette.

The take-home message

QT interval can be abnormally long or short. Although we usually look for an abnormally long QT interval, we should not forget about short QT interval.

References

- 1. Ahmed R, Hashiba K. Reliability of QT intervals as indicators of clinical hypercalcemia. Clin Cardiol 1988;11(6):395-400.
- 2.Schimpf R, Wolpert C, Gaita F, Giustetto C, Borggrefe M. Short QT syndrome. Cardiovasc Res 2005;67(3):357-66.
- 3.Vella A, Gerber TC, Hayes DL, Reeder GS. Digoxin, hypercalcaemia, and cardiac conduction. Postgrad Med J 1999;75(887):554-6.

www.ijmbs.org

Reviewers

NA (solicited submission)

Editors

Salem A Beshyah, Abu Dhabi, UAE. Elmahdi Elkhammas, Columbus, Ohio, USA.