

Utility of hepatitis C virus RNA as the screening test for diagnosing hepatitis C virus infection in hemodialysis patients

Sir,

Hepatitis C virus (HCV) is a major life threat among hemodialysis (HD) patients with prevalence of 4.3%-45.2% in India.^[1] Major challenge is that anti-HCV which is usually a reliable marker for HCV is often negative in HD patients despite the presence of significant viremia, making molecular tests the most reliable tests for screening HCV infection in HD patients.^[2] Goal of this study was to find out the utility of HCV RNA as a screening marker for HCV infection in HD patients.

In this one year retrospective study, blood samples of 410 chronic kidney disease (CKD) patients were included. Anti-HCV was tested by chemiluminescence microparticle immunoassay (Architect i1000SR, Abbott Diagnostics, USA), and HCV RNA was tested by real-time PCR (Abbott m2000rt). Institutional Ethics Committee of our institute approved the study protocol.

Among 410 CKD patients, 128 (31%) were on HD and overall HCV infection among 128 HD patients was seen in 17 (13.3%) patients with 12 (9.4%) patients positive for anti-HCV and 12 (9.4%) patients positive for HCV RNA. A total of 7 (5.5%) patients were positive for both anti-HCV and HCV RNA suggesting active infection. A total of 5 (4%) patients were HCV RNA negative but anti-HCV positive with low signal/cut-off ratio who persistently remained HCV RNA negative till 1 year follow-up, thus suggesting false anti-HCV positivity.^[3] Totally 5 (4%) asymptomatic HD patients with undetectable antibodies were HCV RNA positive proving that anti-HCV alone is not a good screening marker for HCV detection among HD patients table 1. Our study results correlate well with two other studies.^[4,5]

Seronegative HCV infection among HD patients can be due to longer window period in these immunocompromised individuals^[4] or due to the presence of antigen-antibody immune complexes, resulting in low anti-HCV response.^[2] Hence, we recommend that all HD patients should be screened by both anti-HCV and HCV RNA to rule out HCV infection with high sensitivity.

Table 1: Hepatitis C virus viral markers in hemodialysis patients in the study group (n=128)

HCV viral markers in HD patients	Total number (%)
Total anti-HCV positive	12 (9.4)
Total HCV RNA positive	12 (9.4)
Anti-HCV positive and HCV RNA positive	7 (5.5)
Anti-HCV positive and HCV RNA negative	5 (4)
Anti-HCV negative and HCV RNA positive	5 (4)

HD = Hemodialysis, HCV = Hepatitis C virus

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Supriya Mahajan, Suman Lata Nayak¹, Ekta Gupta

Department of Clinical Virology, Institute of Liver and Biliary Sciences, ¹Department of Nephrology, Institute of Liver and Biliary Sciences, New Delhi, India
E-mail: ektagaurisha@gmail.com

References

- Saha D, Agarwal SK. Hepatitis and HIV infection during haemodialysis. *J Indian Med Assoc* 2001;99:194-9.
- Kazmierczak J, Pawelczyk A, Cortes KC, Radkowski M. Seronegative hepatitis C virus infection. *Arch Immunol Ther Exp (Warsz)* 2014;62:145-51.
- Gupta E, Bajpai M, Choudhary A. Hepatitis C virus: Screening, diagnosis, and interpretation of laboratory assays. *Asian J Transfus Sci* 2014;8:19-25.
- Jindal N, Sooin D, Grover P, Bansal R, Malhotra R, Singh S, *et al.* Hepatitis C virus (HCV) infection among seronegative patients undergoing haemodialysis in a remotely located tertiary care hospital of Northern India: Value of HCV-RNA and Genotypes. *J Clin Diagn Res* 2015;9:DC 10-2.
- Moini M, Ziyaeyan M, Aghaei S, Sagheb MM, Taghavi SA, Moeini M, *et al.* Hepatitis C virus (HCV) infection rate among seronegative hemodialysis patients screened by two methods; HCV core antigen and polymerase chain reaction. *Hepat Mon* 2013;13:e9147.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code: 	Website: www.jlponline.org
	DOI: 10.4103/JLP.JLP_99_17

How to cite this article: Mahajan S, Nayak SL, Gupta E. Utility of hepatitis C virus RNA as the screening test for diagnosing hepatitis C virus infection in hemodialysis patients. *J Lab Physicians* 2017;9:345.
© 2017 Journal of Laboratory Physicians | Published by Wolters Kluwer - Medknow