# **Preventing Newborn Deaths in Marginalized Population Groups**

## Dear Editor,

Since the time of adoption of the millennium development goals, a remarkable success has been observed in the field of saving lives of under-five children, including newborns.<sup>[1]</sup> In fact, in excess of a 60% decline has been observed in the incidence of under-five deaths from 1990 to 2016; nevertheless, the progress has not been universal.<sup>[2]</sup> In 2016 alone, more than 45% of reported under-five deaths were in the neonatal age group, which is an alarming cause of concern.<sup>[2]</sup> To attain universal health coverage and promote the survival of newborns, the need of the hour is to target vulnerable sections of the community.<sup>[1,2]</sup>

Further, based on the available estimates, it has been anticipated that 30 million newborns will die in the next 13 years from 2017.<sup>[2]</sup> This should act as the driving force for improving the quality of services and ensuring the provision of timely care during and after childbirth.<sup>[1,2]</sup> The available recent global figures suggest that India accounts for the maximum number of newborn deaths followed by Pakistan, while preterm birth-related complications, pneumonia, and diarrhea were the most common reasons for the deaths.<sup>[2]</sup> These estimates are an indirect reflection of the high incidence of home deliveries, poor infrastructure/logistics supply in health facilities, lack of funding, poor quality of services, and limited number of trained health professionals in healthcare establishments.<sup>[2]</sup>

It is worth noting that if the global stakeholders fail to improve the progress, in excess of 60 nations will not meet the set target of ending preventable newborn deaths by 2030.<sup>[1,2]</sup> Because the lives of more than 50 million under-five children have been saved since the start of the 21<sup>st</sup> century, it is obvious that the health sector has made significant progress in the field of knowledge and technology desired to save lives.<sup>[2,3]</sup> Thus, it is high time that efforts are taken to minimize the number of deaths of newborns; otherwise, all gains achieved till date is incomplete.<sup>[11]</sup> In other words, it is high time that we take measures to ensure that essential services reach to those who are in utmost need of the same.<sup>[11]</sup>

As the majority of the newborn deaths are preventable through simple and cost-effective interventions, the primary objective is to minimize the existing inequity among different sections of people.<sup>[1,3]</sup> In addition, measures such as enhancing access to skilled personnel during the antenatal period and during childbirth, discouraging the practice of home delivery, provision of basic services (such as immunization, drugs, and promotion of breastfeeding), increasing funding for the program, and enhancing access to water and sanitation facilities can be implemented.<sup>[2-4]</sup> These measures will not only improve the survival chances of the newborn but also even the mother and thus will have a double-folded benefit.<sup>[4]</sup>

To conclude, the proportion of newborn deaths among the total under-five deaths has shown a rise despite the significant progress made in the field of childbirth. This essentially advocates for the necessity to expand the services to the marginalized population groups.

### Financial support and sponsorship Nil

## **Conflicts of interest**

There are no conflicts of interest.

#### Saurabh RamBihariLal Shrivastava, Prateek Saurabh Shrivastava<sup>1</sup>

Department of Community Medicine, Member of the Medical Education Unit and Medical Research Unit, Shri Sathya Sai Medical College and Research Institute, <sup>1</sup>Department of Community Medicine, Shri Sathya Sai Medical College and Research Institute, Kancheepuram, Tamil Nadu, India

Address for correspondence: Dr. Saurabh RamBihariLal Shrivastava, 3<sup>rd</sup> Floor, Department of Community Medicine, Shri Sathya Sai Medical College and Research Institute, Ammapettai Village, Thiruporur - Guduvancherry Main Road, Sembakkam Post, Kancheepuram - 603 108, Tamil Nadu, India. E-mail: drshrishri2008@gmail.com

## REFERENCES

- United Nations Children's Fund. Levels and Trends in Child Mortality; 2017. Available from: https://www.data.unicef.org/resources/ levels-trends-child-mortality. [Last accessed on 2018 Nov 08].
- World Health Organization. Newborns: Reducing Mortality-Fact Sheet No. 333; 2017. Available from: http://www.who.int/mediacentre/ factsheets/fs333/en. [Last accessed on 2018 Nov 07].
- Shrivastava SR, Shrivastava PS, Ramasamy J. Standardizing the assessment and management protocol of critically ill under-five children: World Health Organization. Ann Trop Med Public Health 2016;9:215-6.
- Shrivastava SR, Shrivastava PS, Ramasamy J. World Health Organization launches a network to improve quality of care for mothers, newborns and children. Al Ameen J Med Sci 2017;10:184-5.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: journal.limu.edu.ly
	DOI: 10.4103/LIUJ.LIUJ_48_18

How to cite this article: Shrivastava SR, Shrivastava PS. Preventing newborn deaths in marginalized population groups. Libyan Int Med Univ J 2019;4:43. Received: 07-11-2018 Accepted: 22-11-2018 Published: 12-02-2019 © 2019 Libyan International Medical University Journal | Published by Wolters Kluwer - Medknow