

The Orbitomeningeal Band as a Way to Bloodless Transcavernous Dissection and Anterior Clinoidectomy

Abstract

The meningo-orbital band (MOB) is a dural fold which runs along the lateral border of the superior orbital fissure and contains few small dural veins and the orbitomeningeal artery. MOB detachment is relatively easy to understand step-wise procedure, provides a wider exposure, and better orientation thus facilitating relatively easy approach to paraclinoid and cavernous sinus region. The present microsurgical technique helps to preserve the true cavernous membrane and thereby providing almost bloodless dissection of the cavernous sinus. The same technique can be used to uncover the anterior clinoid process laterally, posteriorly, superiorly, and also in the inferolateral region thereby decreasing the risk and time of clinoidectomy.

Keywords: Anterior clinoid process, clinoidectomy, orbitomeningeal band, paraclinoid region

Introduction

Anterior clinoidectomy (AC) is a technique that is of paramount while clipping the aneurysms involving the internal carotid artery (ICA) and resecting the tumors in the paraclinoid region.^[1-3] Due to deep location and proximity to the vital structures of anterior clinoid process (ACP), the AC is a technically challenging surgical procedure.^[4-6] We describe a technique which helps to preserve the true cavernous membrane thereby providing almost bloodless dissection of the cavernous sinus.

Surgical Technique

The orbitomeningeal band is found on the lateral most edge of the superior orbital fissure. It has been common knowledge that there are no cranial nerves on the lateral most border of the superior orbital fissure. Hence, cutting the orbitomeningeal band on the lateral aspect does not produce any cranial nerve deficit. However, we intend to take this technique a step forward. After we cut the orbitomeningeal band, we enter a compartment since the cut orbitomeningeal band separates into two folds. Using the microscope one can dissect between these folds. As one develops the compartment between the folds using sharp dissection inferiorly, the true cavernous sinus membrane is preserved

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

and the cavernous sinus can be uncovered. Developing the compartment by blunt and sharp dissection superiorly will help to uncover the ACP [Figures 1-5]. This technique helps in cases such as skull base tumors, aneurysms, and trauma when one needs to combine cisternal opening with or without a decompressive hemicraniectomy. It helps in avoiding troublesome cavernous sinus bleeding which often requires fibrin glue injection into the cavernous sinus thus producing an iatrogenic lateral cavernous sinus thrombosis.

Discussion

The sharp posterior border of the lesser wing of the sphenoid (the sphenoid ridge) projects into the lateral sulcus and apex which ends medially is the ACP, which gives rise to the attachment to the tentorium cerebelli.^[7-10] ACP is closely related to many vital neurovascular structures which include optic nerves and ICA.^[11] The meningo-orbital band (MOB), orbitotemporal periosteal fold, or frontotemporal fold is a dural fold which runs along the lateral border of the superior orbital fissure and contains small dural veins and the orbitomeningeal artery and encountered during conventional frontotemporal craniotomies.^[4,12,13]

The objective of AC is to remove the bony process and to create the clinoid

How to cite this article: Cherian I, Kasper EM, Agarwal A. The orbitomeningeal band as a way to bloodless transcavernous dissection and anterior clinoidectomy. *Asian J Neurosurg* 2018;13:943-5.

Iype Cherian,
Ekkehard M.
Kasper¹,
Amit Agarwal²

Department of Neurosurgery,
College of Medical sciences,
Bharatpur, Chitwan, Nepal,
¹Department of Neurosurgery,
Harvard Medical School and
Director of Neurosurgical
Oncology at BIDMC, Boston/
MA, USA, ²Department of
Neurosurgery, Narayana
Medical College and Hospital,
Chintareddypalem, Nellore,
Andhra Pradesh, India

Address for correspondence:

Dr. Iype Cherian,
College of Medical Sciences,
Bharatpur, Chitwan, Nepal.
E-mail: drrajucherian@gmail.
com

Access this article online

Website: www.asianjns.org

DOI: 10.4103/ajns.AJNS_198_16

Quick Response Code:






Figure 1: A sequential understanding of the superior orbital fissure anatomy




Figure 2: The shape of the superior orbital fissure with a vertical and lateral horizontal aspect




Figure 3: The location of the cranial nerves passing through the fissure




Figure 4: The cranial nerves within the outer layer of the cavernous sinus and true cavernous membrane (blue color)

space to reduce the retraction on the brain and allows easy access the intended structures.^[14,15] While opening the orbitotemporal periosteal fold exposes the structures which lie in cleavage plane between the inner layer of dura mater (temporal dura propria) and lateral wall of the cavernous sinus (lacrimal, oculomotor, trochlear, ophthalmic [V1], and abducens nerves).^[16,17] The classic Dolenc's approach includes an AC and a transcavernous dissection.^[3] Majority of the experts perform this procedure by drilling of the ACP after exposing it laterally and superiorly and also peel the dura off the cavernous sinus starting at V2.^[1,2]

The extradural ACs have been described to approach the surgical lesions in the paraclinoid space and anterior cavernous sinus without detachment of the MOB,^[4,12,13,18,19] however in contrast to intradural approach, the extradural AC approach requires more experience and a detail understanding of regional anatomy.^[20] MOB detachment is relatively easy to understand step-wise procedure, provides a wider exposure, and better orientation to approach paraclinoid and cavernous sinus region thus facilitates easier manipulation of the lesion and reduces chances of

transmission of direct heat and injury to the surrounding structures.^[4,12-14,21-24]

Conclusion

The present microsurgical technique helps to preserve the true cavernous membrane and thereby providing almost bloodless dissection of the cavernous sinus. The same technique can be used to uncover the ACP laterally, posteriorly, superiorly, and also in the inferolateral region thereby decreasing the risk and time of clinoidectomy.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

1. Mikami T, Minamida Y, Koyanagi I, Baba T, Houkin K. Anatomical variations in pneumatization of the anterior clinoid process. *J Neurosurg* 2007;106:170-4.




Figure 5: The orbitomenigeal artery at the lateral end of the orbitomenigeal band which needs to be coagulated before the dissection starts; the anterior clinoid process

2. Bayassi S. Meningo-orbital fold (MOF) as a guiding point in extradural approach to the anterior clinoid process. *Neurol Neurochir Pol* 2005;39:49-55.
3. Dolenc VV. A combined epi- and subdural direct approach to carotid-ophthalmic artery aneurysms. *J Neurosurg* 1985;62:667-72.
4. Coscarella E, Baskaya MK, Morcos JJ. An alternative extradural exposure to the anterior clinoid process: The superior orbital fissure as a surgical corridor. *Neurosurgery* 2003;53:162-6.
5. Drake CG, Vanderlinden RG, Amacher AL. Carotid-ophthalmic aneurysms. *J Neurosurg* 1968;29:24-31.
6. Hauser MJ, Gass H. Optic nerve pressure by aneurysm relieved by decompression of optic nerve; report of a case. *AMA Arch Ophthalmol* 1952;48:627-31.
7. Inoue T, Rhoton AL Jr., Theele D, Barry ME. Surgical approaches to the cavernous sinus: A microsurgical study. *Neurosurgery* 1990;26:903-32.
8. Evans KL, Crowder J, Miller ES. Subtilisins of *Bacillus* spp. hydrolyze keratin and allow growth on feathers. *Can J Microbiol* 2000;46:1004-11.
9. Sade B, Kweon CY, Evans JJ, Lee JH. Enhanced exposure of carotico-oculomotor triangle following extradural anterior clinoidectomy: A comparative anatomical study. *Skull Base* 2005;15:157-61.
10. Rhoton AL Jr. The anterior and middle cranial base. *Neurosurgery* 2002;51 4 Suppl: S273-302.
11. Romani R, Elsharkawy A, Laakso A, Kangasniemi M, Hernesniemi J. Complications of anterior clinoidectomy through lateral supraorbital approach. *World Neurosurg* 2012;77:698-703.
12. Ammirati M, Bernardo A. Anatomical study of the superior orbital fissure as seen during a pterional approach. *J Neurosurg* 2007;106:151-6.
13. Noguchi A, Balasingam V, Shiokawa Y, McMenomey SO, Delashaw JB Jr. Extradural anterior clinoidectomy. Technical note. *J Neurosurg* 2005;102:945-50.
14. Froelich SC, Aziz KM, Levine NB, Theodosopoulos PV, van Loveren HR, Keller JT. Refinement of the extradural anterior clinoidectomy: Surgical anatomy of the orbitotemporal periosteal fold. *Neurosurgery* 2007;61 5 Suppl 2:179-85.
15. Kim JM, Romano A, Sanan A, van Loveren HR, Keller JT. Microsurgical anatomic features and nomenclature of the paracaloid region. *Neurosurgery* 2000;46:670-80.
16. Lehmeberg J, Krieg SM, Meyer B. Anterior clinoidectomy. *Acta Neurochir (Wien)* 2014;156:415-9.
17. Kapur E, Mehic A. Anatomical variations and morphometric study of the optic strut and the anterior clinoid process. *Bosn J Basic Med Sci* 2012;12:88-93.
18. Day JD, Giannotta SL, Fukushima T. Extradural temporo-polar approach to lesions of the upper basilar artery and infrachiasmatic region. *J Neurosurg* 1994;81:230-5.
19. Yonekawa Y, Ogata N, Imhof HG, Olivecrona M, Strommer K, Kwak TE, *et al*. Selective extradural anterior clinoidectomy for supra- and parasellar processes. Technical note. *J Neurosurg* 1997;87:636-42.
20. Yoon BH, Kim HK, Park MS, Kim SM, Chung SY, Lanzino G. Meningeal layers around anterior clinoid process as a delicate area in extradural anterior clinoidectomy: Anatomical and clinical study. *J Korean Neurosurg Soc* 2012;52:391-5.
21. François P, Travers N, Lescanne E, Arbeille B, Jan M, Velut S. The interperiosteal-dural concept applied to the perisellar compartment: A microanatomical and electron microscopic study. *J Neurosurg* 2010;113:1045-52.
22. Umansky F, Nathan H. The lateral wall of the cavernous sinus. With special reference to the nerves related to it. *J Neurosurg* 1982;56:228-34.
23. Kawase T, van Loveren H, Keller JT, Tew JM. Meningeal architecture of the cavernous sinus: Clinical and surgical implications. *Neurosurgery* 1996;39:527-34.
24. Fukuda H, Evins AI, Burrell JC, Iwasaki K, Stieg PE, Bernardo A. The meningo-orbital band: Microsurgical anatomy and surgical detachment of the membranous structures through a frontotemporal craniotomy with removal of the anterior clinoid process. *J Neurol Surg B Skull Base* 2014;75:125-32.