

Diabetic Peripheral Neuropathy: A Systematic Review of Nigerian Patients

Taoreed Adegoke Azeez, Emmanuel Chinedu Eguzozie, Oladotun Victor Olalusi¹

Department of Medicine, Endocrinology Unit, University College Hospital, ¹Department of Medicine, Neurology Unit, University College Hospital, Ibadan, Nigeria

Abstract

Diabetes mellitus is associated with microvascular and macrovascular complications. Diabetic neuropathy is the most common long-term complication of diabetes. The study was aimed at determining the prevalence of diabetic peripheral neuropathy and identifying its associated factors. Medical databases, including PubMed, Google Scholar, African Journal Online, and SCOPUS were searched and eligible studies were selected using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses algorithm. The prevalence of diabetic peripheral neuropathy in Nigeria was dependent on the modality used in diagnosing peripheral neuropathy. The overall prevalence of diabetic peripheral neuropathy in Nigeria was 31.2%–97.5%. The modality-dependent prevalences were 37%–97.5% (by biothesiometry), 41.7%–75% (by Michigan Neuropathy Screening Instrument), 31.2%–43.3% (by United Kingdom screening test), and 43.3%–69.9% (by diabetic neuropathic examination score). The associated factors were duration and control of diabetes, the age of the patient, presence of cardiovascular risk factors such as hypertension and dyslipidemia and other microvascular complications (e.g., diabetic retinopathy and nephropathy). The prevalence of diabetic peripheral neuropathy was found to be high in Nigeria and the associated risk factors were age, glycemic control, and cardiovascular risk factors.

Keywords: Diabetic neuropathy, diagnosis, Nigeria, prevalence, risk factors

INTRODUCTION AND GENERAL DEFINITIONS

Diabetes mellitus comprises a group of heterogeneous metabolic disorders in which the common denominator is hyperglycemia resulting from a defect in insulin secretion, action, or both.^[1] The prevalence is rising rapidly across the various continents of the world.^[2] The estimated prevalence of diabetes mellitus in Africa is 3.1%.^[3] In Sub-Saharan Africa, Nigeria has the highest number of persons living with diabetes mellitus.^[4] According to the International Diabetes Federation, as of 2020, about 2.7 million adults had diabetes in Nigeria.^[5] Figure 1 shows the regional distribution of diabetes mellitus prevalence rates across the various geopolitical zones of Nigeria.^[6]

DIABETIC MICROANGIOPATHY

Chronic hyperglycemia induces pathological changes in the microvasculature of nerves, retina, and glomeruli resulting in the development of diabetic microangiopathy.^[7] These are the underlying mechanisms for the development of diabetic neuropathy, retinopathy, and nephropathy. There are

various pathways involved in the development of diabetic microangiopathy. Hyperglycemia, through the advanced glycation end products, polyol and hexosamine pathways, triggers an injurious cascade to the basement membrane of the microvasculature.^[8] Hyperosmotic stress can induce inflammation by stimulating the release of pro-inflammatory cytokines, which has been found to contribute to diabetic microangiopathy.^[9] The role of oxidative stress in the development and progression of microangiopathy has been extensively described in the literature.^[10] Endothelial dysfunction, through deranged nitric oxide synthase function, has also been documented to contribute to diabetic microangiopathy.^[11] The mechanisms underlying diabetic microangiopathy are summarized in Figure 2.

Address for correspondence: Dr. Taoreed Adegoke Azeez,
Department of Medicine, Endocrinology Unit, University College Hospital,
Ibadan, Nigeria.
E-mail: adegokegalaxy@yahoo.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Azeez TA, Eguzozie EC, Olalusi OV. Diabetic peripheral neuropathy: A systematic review of Nigerian patients. *Libyan Int Med Univ J* 2021;6:12-8.

Received: 15-03-2021

Revised: 30-04-2021

Accepted: 23-05-2021

Published: 21-06-2021

Access this article online

Quick Response Code:



Website:
journal.lim.u.edu.ly

DOI:
10.4103/liuj.liuj_62_21

DIABETIC NEUROPATHY

Diabetic neuropathy refers to symptoms and/or signs of nerve dysfunction in a patient with diabetes when other causes of nerve dysfunction have been excluded from the study.^[12] Diabetic neuropathy is the most common long-term complication of diabetes with a huge contribution to the morbidity and mortality of patients with diabetes.^[13] It is also the most common cause of neuropathy, especially in the advanced economies.^[14] In Northern America and Europe, the estimated prevalence of diabetic neuropathy is 6%–51% depending on the population that was studied.^[15] In a meta-analysis involving 23 studies done across Africa, the estimated prevalence of diabetic neuropathy in Africa was 46%.^[16] In another meta-analysis, the identified factors associated with diabetic neuropathy were duration of diabetes, age, glycated hemoglobin, and diabetic retinopathy.^[17] Table 1 shows the classification of diabetic neuropathy, as published by the American Diabetes Association.^[18] Diabetic symmetric polyneuropathy is the most common form of diabetic neuropathy.^[19]

The pathophysiology of diabetic peripheral neuropathy involves both metabolic and vascular mechanisms.^[20] As evidenced by nerve biopsy studies, the nerve microvasculature showed thickening of the basement membrane as well as hyperplasia and hypertrophy of endothelial cells.^[21] The metabolic pathways usually involve reduction-oxidation stress.^[22] Diabetic peripheral neuropathy often starts from the toes until it involves both the lower and upper limbs in the gloves and stockings distribution pattern.^[23] A significant proportion of patients with diabetic peripheral neuropathy present with painful symptoms such as dysaesthesia, allodynia, and hyperaesthesia.^[24] Sometimes, the patient presents with numbness and it is not uncommon for patients with diabetes to be asymptomatic of diabetic peripheral neuropathy.^[25] The various modalities for diagnosing

diabetic peripheral neuropathy include nerve conduction studies (the gold standard), nerve biopsy, various scoring methods using signs and symptoms of peripheral neuropathy (such as Michigan Neuropathy Screening Instrument, Diabetic Neuropathy Symptoms Score, and Toronto Clinical Scoring System), vibration threshold potential using a biothesiometer, and using a monofilament.^[25] Based on the reports of previous studies, the sensitivity of the various modalities of diagnosing diabetes ranged from 63% to 98%.^[26-28]

Objectives

1. To determine the prevalence of diabetic peripheral neuropathy in Nigeria
2. To identify the factors associated with the development of prevalence of diabetic peripheral neuropathy in Nigeria.

METHODS

Medical databases including African Journals online, Google Scholar, Medline, and SCOPUS were used to retrieve studies on diabetic peripheral neuropathy in Nigeria. The search terms used were “diabetic neuropathy,” “diabetic peripheral neuropathy,” “peripheral neuropathy,” and “diabetic symmetrical polyneuropathy.” Other terms used in the data search included “diabetic neuropathy in Nigeria,” “diabetic microangiopathy,” “complications of diabetes” and “diabetic peripheral neuropathy in Nigeria.” In order to improve the quantity and quality of retrieved articles, Boolean operators such as “AND” as well as “OR” were used. The gray literature was also searched to enhance the depth of the retrieved studies. The literature search was done by strictly following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. The PRISMA flow diagram is shown in Figure 3.

Inclusion criteria

1. Studies done/published between January 1, 2000, and December 31, 2020, to determine the prevalence of diabetic peripheral neuropathy and its associated factors in Nigeria




Figure 1: Prevalence of diabetes mellitus across the geopolitical zones in Nigeria

Table 1: Classification of diabetic neuropathy

Types of diabetic neuropathy
Generalized
Distal symmetric polyneuropathy (typical and atypical)
Focal/multifocal
Mononeuropathy
Multiple mononeuropathies
Lumbosacral, thoracic, and cervical radiculoplexus neuropathies
Autonomic neuropathy




Figure 2: Pathophysiology of diabetic microangiopathy




Figure 3: The PRISMA flow diagram of the literature search and selection

2. Studies whose abstracts and or full text were available at the searched databases or from the gray literature.

Exclusion criteria

1. Studies on diabetic peripheral neuropathy outside Nigeria
2. Studies on diabetic peripheral neuropathy inside Nigeria but not focusing on the prevalence of diabetic peripheral neuropathy or its associated risk factors in Nigeria.

The studies were identified and screened independently by the authors. The ones selected were agreed upon by at least two out of the three authors. Data extraction was done on a spreadsheet and the results were presented as tables and charts.

RESULTS

Thirteen studies were selected for the systematic review. The range of the average age in the systematic review was 42–60 years. The various studies and the geopolitical zones are shown in Table 2. All the selected studies were cross-sectional studies. The sample size in each study is shown in Table 3. The total sample size was 2848.

The various modalities used in diagnosing diabetic peripheral neuropathy are shown in Table 4. Some studies used two different modalities in diagnosing diabetic peripheral neuropathy. The four modalities of diagnosing diabetic peripheral neuropathy employed in the various studies include biothesiometry, Michigan Neuropathy Screening Instrument, United Kingdom Screening Test, and diabetic neuropathy examination score. The most commonly used modality was biothesiometry.

The prevalence of diabetic peripheral neuropathy in each study is shown in Table 5. The prevalence of diabetic peripheral neuropathy in this systematic review was 31.2%–97.5%. The prevalence of diabetic peripheral neuropathy with respect to the modality of diagnosis is shown in Table 6. Averagely, the prevalence of diabetic peripheral neuropathy in Nigeria is lowest using the United Kingdom screening test and highest using the biothesiometry modality.

The associated risk factors for diabetic peripheral neuropathy according to the different studies are shown in As shown in Figure 4, the most commonly reported risk factors for diabetic peripheral neuropathy in Nigeria were the duration of diabetes and the current age of the patient as of the time of performing each study.

DISCUSSION

The frequency of diabetic peripheral neuropathy in Nigeria,

Table 2: List of selected studies across the geopolitical zones

Geopolitical zone	Studies	Year of publication
South-West	Owolabi and Ipadeola ^[29]	2012
	Olamoyegun <i>et al.</i> ^[30]	2015
	Ogbera <i>et al.</i> ^[31]	2015
	Adesina <i>et al.</i> ^[32]	2016
North-Central	Ugoya <i>et al.</i> ^[33]	2006
	Uwakwe <i>et al.</i> ^[34]	2017
	Bello <i>et al.</i> ^[35]	2019
South-East	Asomugha ^[36]	2012
	Ede <i>et al.</i> ^[37]	2018
North-West	Ibrahim <i>et al.</i> ^[38]	2015
	Kaoje <i>et al.</i> ^[39]	2020
North-East	Salawu <i>et al.</i> ^[40]	2018
Multicentric, involving many zones	Chinenye <i>et al.</i> ^[41]	2012

Table 3: Sample size and prevalence rates in the selected studies

Study	Sample size	Prevalence (%)
Adesina <i>et al.</i> ^[32]	50	50.0
Olamoyegun <i>et al.</i> ^[30]	90	69.6
Uwakwe <i>et al.</i> ^[34]	100	75
Ede <i>et al.</i> ^[37]	100	73.3
Ugoya <i>et al.</i> ^[33]	120	75
Bello <i>et al.</i> ^[35]	175	41.7
Asomugha ^[36]	200	97.5
Ogbera <i>et al.</i> ^[31]	225	37
Salawu <i>et al.</i> ^[40]	250	86.8
Owolabi and Ipadeola ^[29]	277	71.1
Kaoje <i>et al.</i> ^[39]	330	39.7
Ibrahim <i>et al.</i> ^[38]	400	31.2
Chinenye <i>et al.</i> ^[41]	531	59.2
Total	2848	

Table 4: Modalities for diagnosing diabetic peripheral neuropathy in various studies

Modality of diagnosis	Studies
Biothesiometry using VPT	Salawu <i>et al.</i> ^[40] Asomugha ^[36] Kaoje <i>et al.</i> ^[39] Uwakwe <i>et al.</i> ^[34] Olamoyegun <i>et al.</i> ^[30] Adesina <i>et al.</i> ^[32] Ogbera <i>et al.</i> ^[31]
United Kingdom screening test	Ibrahim <i>et al.</i> ^[38] Kaoje <i>et al.</i> ^[39]
Diabetic neuropathic examination score	Asomugha ^[36] Olamoyegun <i>et al.</i> ^[30]
Michigan neuropathy screening instrument	Bello <i>et al.</i> ^[35] Ede <i>et al.</i> ^[37] Ugoya <i>et al.</i> ^[33] Owolabi and Ipadeola ^[29]
VPT: Vibration perception threshold	

according to this systematic review, was 37.2%–97.5%. This is a wide range largely due to differences in sociodemographics and methodology. In a systematic review and meta-analysis, involving studies across the African continent, the prevalence rate of diabetic peripheral neuropathy ranged between 16.6% and 83.4%.^[29] The range was also wide and the rate is comparable with the present study. The main reasons put forward for the wide range were differences in sociodemographics, type of diabetes and the methods employed in diagnosing diabetic peripheral neuropathy, just as was found in this systematic review. In another systematic review done in China, the prevalence rate of diabetic peripheral neuropathy was also found to be very wide and variations in demographics and methods of diagnosing the disease were reported as being responsible for this wide range.^[42]

Similarly, in a systematic review done in Iran, the prevalence of diabetic peripheral neuropathy in the various studies ranged from 27.4% to 87.3%. Just as it was stated for the African and Chinese systematic reviews, the authors in the Iranian study also acknowledged a wide range in the prevalence of diabetic peripheral neuropathy and ascribed this wide variation to the differences in sociodemographics and research methods.^[12] The sensitivity and specificity of the various diagnostic modalities for diabetic peripheral neuropathy vary, so it is expected that the different methods will yield different prevalence rates.^[12]

The most prominent associated risk factor for diabetic peripheral neuropathy in this study was the duration of diabetes. A similar finding has been reported from other previous systematic reviews in Africa and Asia.^[16,42,43] This is important because the prevalence of early-onset diabetes is increasing in Nigeria due to the adoption of Western lifestyles and the risk of developing peripheral neuropathy with its attendant complication of diabetic foot ulcer will also increase in this population.^[44] It is therefore of crucial importance to prevent or delay the onset of diabetes.

The age of the patients was also another strong predictor of the occurrence of diabetic peripheral neuropathy, according to

Table 5: Prevalence of diabetic peripheral neuropathy using different modalities of diagnosis

Diagnostic modality	Prevalence (%)
Biothesiometry using VPT	37-97.5
Michigan neuropathy screening instrument	41.7-75
United Kingdom screening test	31.2-43.3
Diabetic neuropathic examination score	43.3-69.6

VPT: Vibration perception threshold

the findings of this study. Other studies have also documented age as a prominent factor associated with the development of diabetic peripheral neuropathy.^[17,45,46] It has been reported that nerve conduction velocity declines with age and this may be responsible for the effect of age on the incidence of peripheral neuropathy in individuals with diabetes.^[47]

The present systematic review also showed that poor glycaemic control was associated with diabetic peripheral neuropathy. Similarly, Aleidan *et al.*^[48] in Saudi Arabia, Awadalla *et al.*^[49] in Sudan and Gill *et al.*^[50] in Ethiopia independently reported that poor glycaemic control was associated with diabetic peripheral neuropathy. Studies have shown that prolonged exposure of nerves to hyperglycemia is critically important for the development of diabetic peripheral neuropathy.^[51,52]

This study also found the universal cardiovascular risk factors such as hypertension, obesity, dyslipidaemia, smoking, and alcohol ingestion to be associated with diabetic neuropathy. Smith *et al.*, also reported an association between dyslipidemia as well as obesity and diabetic peripheral neuropathy.^[53] Papanas and Ziegler, also reported a strong association between




Figure 4: Risk factors for diabetic peripheral neuropathy and the number of studies reporting each

Table 6: Risk factors for diabetic peripheral neuropathy

Risk factors	Studies
Current age of the patient	Salawu <i>et al.</i> ^[40] Asomugha ^[36] Bello <i>et al.</i> ^[35] Ede <i>et al.</i> ^[37] Uwakwe <i>et al.</i> ^[34] Olamoyegun <i>et al.</i> ^[30] Ogbera <i>et al.</i> ^[31] Owolabi and Ipadeola ^[29]
Diabetes duration	Salawu <i>et al.</i> ^[40] Ibrahim <i>et al.</i> ^[38] Asomugha ^[36] Bello <i>et al.</i> ^[35] Ede <i>et al.</i> ^[37] Uwakwe <i>et al.</i> ^[34] Olamoyegun <i>et al.</i> ^[30] Adesina <i>et al.</i> ^[32]
Male gender	Asomugha ^[36] Olamoyegun <i>et al.</i> ^[30]
Poor glycemic control	Bello <i>et al.</i> ^[35] Ede <i>et al.</i> ^[37] Olamoyegun <i>et al.</i> ^[30] Ogbera <i>et al.</i> ^[31]
Elevated blood pressure	Salawu <i>et al.</i> ^[40] Bello <i>et al.</i> ^[35] Olamoyegun <i>et al.</i> ^[30]
Age at the time of diagnosing diabetes	Ibrahim <i>et al.</i> ^[38]
Diabetic retinopathy	Ibrahim <i>et al.</i> ^[38]
Proteinuria	Ibrahim <i>et al.</i> ^[38]
Smoking	Ede <i>et al.</i> ^[37]
Alcohol ingestion	Ede <i>et al.</i> ^[37]
Body mass index	Olamoyegun <i>et al.</i> ^[30]
Dyslipidemia	Owolabi and Ipadeola ^[29]

traditional cardiovascular risk factors such as hypertension, obesity, dyslipidaemia, smoking and alcohol ingestion, and diabetic peripheral neuropathy.^[54]

In addition, the presence of other microangiopathic disorders such as diabetic retinopathy and diabetic nephropathy was found to be associated with the development of diabetic peripheral neuropathy. Other authors have documented a similar association among the microvascular complications of diabetes.^[55-57]

CONCLUSION

Diabetic peripheral neuropathy is a common microvascular complication among patients with diabetes mellitus in Nigeria. The factors associated with the presence of diabetic peripheral neuropathy are the duration of diabetes, age of the patient, and cardiovascular risk factors such as hypertension and dyslipidemia and other microvascular complications such as diabetic retinopathy and nephropathy.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Harreiter J, Roden M. Diabetes mellitus-Definition, classification, diagnosis, screening and prevention (Update 2019). *Wien Klin Wochenschr* 2019;131:6-15.
- Harding JL, Pavkov ME, Magliano DJ, Shaw JE, Gregg EW. Global trends in diabetes complications: A review of current evidence. *Diabetologia* 2019;62:3-16.
- Kibirige D, Lumu W, Jones AG, Smeeth L, Hattersley AT, Nyirenda MJ. Understanding the manifestation of diabetes in sub Saharan Africa to inform therapeutic approaches and preventive strategies: A narrative review. *Clin Diabetes Endocrinol* 2019;5:2.
- Dahiru T, Aliyu AA, Sheu AU. A review of population-based studies of diabetes in Nigeria. *Sub Saharan Afr J Med* 2016;3:59-64.
- International Diabetes Federation. IDF Africa Members. Available from: <https://idf.org/our-network/regions-members/africa/members/20-nigeria.html>. [Last accessed on 2021 Jan 20].
- Uloko AE, Musa BM, Ramalan MA, Gezawa ID, Puepet FH, Uloko AT,

et al. Prevalence and risk factors for diabetes mellitus in Nigeria: A systematic review and meta-analysis. *Diabetes Ther* 2018;9:1307-16.

- Chawla A, Chawla R, Jaggi S. Microvascular and macrovascular complications in diabetes: Distinct or continuum. *Indian J Endocrinol Metab* 2016;20:546-51.
- Madonna R, Balistreri CR, Geng YJ, De Caterina R. Diabetic microangiopathy: Pathogenetic insights and novel therapeutic approaches. *Vascul Pharmacol* 2017;90:1-7.
- Brocker C, Thompson DC, Vasiliou V. The role of hyperosmotic stress in inflammation and disease. *Biomol Concepts* 2012;3:345-64.
- Steven S, Frenis K, Oelze M, Kalinovic S, Kuntic M, Jimenez MT, *et al.* Vascular inflammation and oxidative stress: Major triggers for cardiovascular diseases. *Oxid Med Cell Longev* 2019;2019:7092151.
- Sena CM, Pereira AM, Seica R. Endothelial dysfunction – A major mediator of diabetic vascular disease. *Biochim Biophys Acta* 2013;1832:2216-31.
- Bansal V, Kalita J, Misra UK. Diabetic neuropathy. *BMJ Postgrad Med J* 2006;82:95.
- Vinik AI, Nevoret ML, Casellini C, Parson H. Diabetic neuropathy. *Endocrinol Metab Clin North Am* 2013;42:747-87.
- Said G. Diabetic neuropathy – A review. *Nat Clin Pract Neurol* 2007;3:331-40.
- Hicks CW, Selvin E. Epidemiology of peripheral neuropathy and lower extremity disease in diabetes. *Curr Diab Rep* 2019;19:86.
- Shiferaw WS, Akalu TY, Work Y, Aynalem YA. Prevalence of diabetic peripheral neuropathy in Africa: A systematic review and meta-analysis. *BMC Endocr Disord* 2020;20:49.
- Liu X, Xu Y, An M, Zeng Q. The risk factors for diabetic peripheral neuropathy: A meta-analysis. *PLoS One* 2019;14:e0212574.
- Tesfaye S, Boulton AJ, Dyck PJ, Freeman R, Horowitz M, Kempner P, *et al.* Diabetic neuropathies: Update on definition, diagnostic criteria, estimation of severity and treatments. *Diabetes Care* 2020;33:2285-93.
- Llewelyn JG. The diabetic neuropathies: Types, diagnosis and management. *J Neurol Neurosurg Psychiatry* 2003;74 Suppl 2:i15-9.
- Tesfaye S, Selvarajah D. Advances in the epidemiology, pathogenesis and management of diabetic peripheral neuropathy. *Diabetes Metab Res Rev* 2012;28 Suppl 1:8-14.
- Malik RA, Tesfaye S, Newrick PG, Walker D, Rajbhandari SM, Siddique I, *et al.* Sural nerve pathology in diabetic patients with minimal but progressive neuropathy. *Diabetologia* 2005;48:578-85.
- Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res* 2010;107:1058-70.
- Marchettini P, Lacerenza M, Mauri E, Marangoni C. Painful peripheral neuropathies. *Curr Neuropharmacol* 2006;4:175-81.
- Vinik A, Cassellini C, Nevoret ML. Diabetic Neuropathies. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK279175/>. [Last accessed on 2021 Jan 20].
- Russell JW, Zilliox LA. Diabetic neuropathies. *Continuum (Minneapolis)* 2014;20:1226-40.

26. Martin CL, Waberski BH, Pop-Busui R, Cleary PA, Catton S, Albers JW, *et al.* Vibration perception threshold as a measure of distal symmetric neuropathy in type 1 diabetes – Results from the DCCT/EDIC study. *Diabetes Care* 2010;33:2635-41.
27. Fateh HR, Madani SP, Heshmat R, Larijani B. Correlation of Michigan neuropathy screening instrument, United Kingdom screening test and electrodiagnosis for early detection of diabetic peripheral neuropathy. *J Diabetes Metab Disord* 2015;15:8.
28. Mythili A, Kumar KD, Subrahmanyam KA, Venkateswarlu K, Butchi RG. A Comparative study of examination scores and quantitative sensory testing in diagnosis of diabetic polyneuropathy. *Int J Diabetes Dev Ctries* 2010;30:43-8.
29. Owolabi MO, Ipadeola A. Total vascular risk as a strong correlate of severity of diabetic peripheral neuropathy in Nigerian Africans. *Ethn Dis* 2012;22:106-12.
30. Olamoyegun M, Ibraheem W, Iwuala S, Audu M, Kolawole B. Burdens and patterns of microvascular complications in type 2 diabetes in a tertiary health institution in Nigeria. *Afr Health Sci* 2015;15:1136-41.
31. Ogbera AO, Adeleye O, Solagberu B, Azenabor A. Screening for peripheral neuropathy and peripheral arterial disease in persons with diabetes mellitus in a Nigerian University Teaching Hospital. *BMC Res Notes* 2015;8:533.
32. Adesina O, Adewusi F, Alalade B, Otukoya O, Olukunle G, Bello A. Correlation between the duration of diabetes mellitus and neuropathy in a Nigerian tertiary hospital. *Endocr Abstr* 2016;44:104.
33. Ugoya SO, Echejoh GO, Ugoya TA, Agaba EI, Puepet FH, Ogunniyi A. Clinically diagnosed diabetic neuropathy: Frequency, types and severity. *J Natl Med Assoc* 2006;98:1763-6.
34. Uwakwe JN, Odoh G, Edah JO, Enanimo M, Chuhwak EK, Puepet FH. Diabetic peripheral neuropathy and its risk factors: A community-based study. *Int Res J Med Biomed Sci* 2017;2:9-13.
35. Bello A, Biliaminu S, Wahab K, Sanya E. Distal symmetrical polyneuropathy and cardiovascular autonomic neuropathy among diabetic patients in Ilorin: Prevalence and predictors. *Niger Postgrad Med J* 2019;26:123-8.
36. Asomugha AL. Assessment of Diabetic Peripheral Neuropathy in Nigerians Using the Diabetic Neuropathy Examination Instrument and Quantitative Sensory Testing – A Fellowship Dissertation Submitted to the National Postgraduate Medical College of Nigeria. Unpublished. Available from: <https://C:/Users/USER/Downloads/628-Article%20Text-3643-1-10-20190326.pdf>. [Last accessed on 2021 Jan 20].
37. Ede O, Eyichukwu GO, Madu KA, Ogbonnaya IS, Okoro KA, Basil-Nwachuku C, *et al.* Evaluation of peripheral neuropathy in diabetic adults with and without foot ulcers in an African population. *J Biomed Sci* 2018;6:71-8.
38. Ibrahim A, Owolabi LF, Borodo MM, Oguniye A. Clinical profile of diabetic sensorimotor polyneuropathy in a tertiary hospital in Northwestern Nigeria. *Niger J Basic Clin Sci* 2015;12:13-9.
39. Kaoje YS, Bello F, Bansi IK, Bakari AG, Yakubu IM, Lawal AY. Prevalence and severity of distal symmetrical polyneuropathy among patients with type 2 diabetes mellitus in Zaria, Nigeria. *Open Access Text* 2020;6:1-4. [doi: 10.15761].
40. Salawu F, Shadrach L, Adenle T, Martins O, Bukbuk D. Diabetic peripheral neuropathy and its risk factors in a Nigerian population with type 2 diabetes mellitus. *Afr J Diabetes Med* 2018;26:16-20.
41. Chinenye S, Uloko AE, Ogbera AO, Ofoegbu EN, Fasanmade OA, Fasanmade AA, *et al.* Profile of Nigerians with diabetes mellitus- Diabcare Nigeria study group (2008): Results of a multicentre study. *Indian J Endocrinol Metab* 2012;16:558-64.
42. Sun J, Wang Y, Zhang X, Zhu S, He H. Prevalence of peripheral neuropathy in patients with diabetes: A systematic review and meta-analysis. *Prim Care Diabetes* 2020;14:435-44.
43. Sobhani S, Asayesh H, Sharifi F, Djalalinia S, Baradaran HR, Arzaghi SM, *et al.* Prevalence of diabetic peripheral neuropathy in Iran: A systematic review and meta-analysis. *J Diabetes Metab Disord* 2014;13:97.
44. Balogun WO, Onasanya AS, Azeez TA. Prevalence and clinical characteristics of Nigerian patients with early-onset type 2 diabetes mellitus. *Niger J Med* 2020;29:49-54.
45. Al-Mahroos F, Al-Roomi K. Diabetic neuropathy, foot ulceration, peripheral vascular disease and potential risk factors among patients with diabetes in Bahrain: A nationwide primary care diabetes clinic-based study. *Ann Saudi Med* 2007;27:25-31.
46. Kiani J, Moghimbeigi A, Azizkhani H, Kosarifard S. The prevalence and associated risk factors of peripheral diabetic neuropathy in Hamedan, Iran. *Arch Iran Med* 2013;16:17-9.
47. Mao F, Zhu X, Liu S, Qiao X, Zheng H, Lu B, *et al.* Age as an independent risk factor for diabetic peripheral neuropathy in chinese patients with type 2 diabetes. *Aging Dis* 2019;10:592-600.
48. Aleidan FA, Ahmad BA, Alotaibi FA, Aleesa DH, Alhefdhi NA, Badri M. Prevalence and risk factors for diabetic peripheral neuropathy among Saudi hospitalized patients: A nested case-control study. *Int J Gen Med* 2020;2020:881-9.
49. Awadalla H, Noor SK, Elmadhoun WM, Almobarak AO, Elmak NE, Abdelaziz SI, *et al.* Diabetes complications in Sudanese individuals with type 2 diabetes: Overlooked problems in sub-Saharan Africa? *Diabetes Metab Syndr* 2017;11 Suppl 2:S1047-51.
50. Gill GV, Gebrekidan A, English P, Wile D, Tesfaye F. Diabetic complications and glycaemic control in remote North Africa. *QJM* 2008;101:793-8.
51. Bansal D, Gudala K, Muthyala H, Esam HP, Nayakallu R, Bhansali A. Prevalence and risk factors of development of peripheral diabetic neuropathy in type 2 diabetes mellitus in a tertiary care setting. *J Diabetes Investig* 2014;5:714-21.
52. Manschot SM, Biessels GJ, Rutten GE, Kessels RP, Gispen WH, Kappelle LJ, *et al.* Peripheral and central neurologic complications in type 2 diabetes mellitus: No association in individual patients. *J Neurol Sci* 2008;264:157-62.
53. Smith AG, Singleton JR. Obesity and hyperlipidaemia are risk factors for early diabetic neuropathy. *J Diabet Complications* 2013;27:436-42.
54. Papanas N, Ziegler D. Risk factors and comorbidities in diabetic neuropathy: An update 2015. *Rev Diabet Stud* 2015;12:48-62.
55. Chandy A, Pawar B, John M, Isaac R. Association between diabetic nephropathy and other diabetic microvascular and macrovascular complications. *Saudi J Kidney Dis Transpl* 2008;19:924-8.
56. Abdollahi A, Moghimi S, Tabasi A, Rajabi MT, Sabet B. Neuropathy and retinopathy in diabetes: Is there any association? *Int J Ophthalmol* 2009;2:57-60.
57. Li J, Cao Y, Liu W, Wang Q, Qian Y, Lu P. Correlations among diabetic microvascular complications: A systematic review and meta-analysis. *Sci Rep* 2019;9:3137.

ملخص المقال باللغة العربية

الاعتلال العصبي المحيطي السكري: مراجعة منهجية للمرضى النيجيريين

المؤلفون:

Taoreed Adegoke AZEEZ, Emmanuel Chinedu EGUZOZIE, Oladotun Victor OLALUSI. Department of Medicine, University College Hospital, Ibadan, Nigeria.

المؤلف المسؤول: Taoreed Adegoke AZEEZ. البريد الإلكتروني: adegokegalaxy@yahoo.com

يرتبط مرض السكري بمضاعفات الأوعية الدموية الدقيقة والكبيرة. يعد الاعتلال العصبي المحيطي السكري من أكثر المضاعفات شيوعاً على المدى الطويل لمرضى السكري. هدفت هذه الدراسة إلى تحديد مدى انتشار الاعتلال العصبي المحيطي السكري وتحديد العوامل المرتبطة به لمرضى نيجيريين. تم البحث في قواعد البيانات الطبية، بما في ذلك (بوب ميد، مانحة جوجل، والمجلات الأفريقية على النت، وكذلك باحث سكوباس)، وتم اختيار الدراسات المؤهلة باستخدام عناصر التقارير المفضلة للمراجعات المنهجية وخوارزمية التحليلات الوصفية (PRISMA) برزما. ولقد وجد أن نسبة تشخيص انتشار الاعتلال العصبي المحيطي السكري في نيجيريا يعتمد بشكل رئيسي على الطريقة المستخدمة في التشخيص. كان معدل انتشار الاعتلال العصبي المحيطي السكري في نيجيريا بصفة عامة يتراوح ما بين 31.2% إلى 97.5%. كانت معدلات الانتشار المعتمدة على التشخيص بطريقة القياس الحيوي تتراوح ما بين 37% إلى 97.5%. أما طريقة التشخيص بواسطة أداة فحص اعتلال الأعصاب لميشيغان فتراوحت ما بين 41.7% إلى 75%. ومن ناحية أخرى فإن طريقة التشخيص بواسطة اختبار فحص المملكة المتحدة تتراوح ما بين 31.2% إلى 43.3%. وأخيراً فإن طريقة التشخيص بواسطة نتيجة فحص مرض السكري العصبي فأعطت نتائج تراوحت ما بين 43.3% إلى 69.9%. كانت العوامل المرتبطة بمعدل انتشار الاعتلال العصبي المحيطي السكري في نيجيريا تشمل مدة مرض السكري والتحكم فيه، وعمر المريض، ووجود عوامل الخطر القلبية الوعائية (مثل ارتفاع ضغط الدم، وخلل الدهون في الدم، ومضاعفات الأوعية الدموية الدقيقة الأخرى مثل اعتلال الشبكية السكري واعتلال الكلية).

الاستنتاج: وجد أن معدل انتشار الاعتلال العصبي المحيطي السكري مرتفع في نيجيريا، وعوامل الخطر المرتبطة بها هي العمر والتحكم في نسبة السكر في الدم وعوامل الخطر القلبية الوعائية.

الكلمات المفتاحية: اعتلال الأعصاب السكري، الانتشار السكري، عوامل الخطر السكري، التشخيص السكري، نيجيريا.