

## Case Report

# Fluorine-18 fluorodeoxyglucose-positron emission tomography/computer tomography in staging of renal cell carcinoma arising from a native kidney with liver and bone metastasis in a renal transplant patient

## ABSTRACT

Renal cell carcinoma (RCC) of the native kidney accounts for <5% of all malignancies found in transplant recipients. There have been only a few reported cases comprising of few renal transplant patients with RCC of native kidneys due to the relative rarity of the condition. Fluorine-18 Fluorodeoxyglucose (FDG)-positron emission tomography (PET)/computed tomography (CT) is used in the staging of RCC. Prognosis of metastatic RCC is poor. We report the first case of 55-year-old postrenal transplant recipient diagnosed with RCC of the native kidney with liver and bone metastases imaged using F-18 FDG PET/CT.

**Keywords:** Fluorine-18 Fluorodeoxyglucose-positron emission tomography/computed tomography, native kidney, renal cell carcinoma, renal transplant

## INTRODUCTION

Renal transplantation remains the best treatment option for patients with end-stage renal disease for their survival and quality of life. There is a significantly increased risk of malignancy in renal transplant recipients as a result of the use of immunosuppressive medication. On the other hand, renal cell carcinoma (RCC) of the native kidney accounts for <5% of all malignancies found in transplant recipients.<sup>[1]</sup> Fluorine-18 Fluorodeoxyglucose (FDG)-positron emission tomography (PET)/computed tomography (CT) is the imaging tool used in the staging of RCC. Our case describes the image findings of F-18 FDG PET/CT in a case of native kidney RCC with liver and bone metastasis in a renal transplant recipient.

## CASE REPORT

A 55-year-old male, postrenal transplant done 5 years before, presented with back pain. ultrasound abdomen was done which showed a hypoechoic lesion in the right native kidney,

suspicious for malignancy. He was referred for the whole body F-18 FDG PET/CT [Figure 1] for staging which showed a cystic lesion in the native right kidney (with a SUVmax of 10, arrows), hypodense liver lesions and multiple skeletal lesions (SUVmax of 15, arrows). He underwent renal biopsy which showed clear cell carcinoma and started on tyrosine kinase inhibitor and is on follow-up.

**KORAMADAI KARUPPUSAMY KAMALESHWARAN, BHARAT RANGARAJAN<sup>1</sup>, RAGHI PARAMBEN JOSE, AJIT SUGUNAN SHINTO**


Department of Nuclear Medicine, PET/CT and Radionuclide Therapy, <sup>1</sup> Department of Medical Oncology, Kovai Medical Center and Hospital Limited, Coimbatore, Tamil Nadu, India

**Address for correspondence:** Dr. Koramadai Karuppusamy Kamaleshwaran, Department of Nuclear Medicine, PET/CT and Radionuclide Therapy, Comprehensive Cancer Care Center, Kovai Medical Center and Hospital Limited, Coimbatore - 641 014, Tamil Nadu, India.  
E-mail: dr.kamaleshwar@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**How to cite this article:** Kamaleshwaran KK, Rangarajan B, Jose RP, Shinto AS. Fluorine-18 fluorodeoxyglucose-positron emission tomography/computer tomography in staging of renal cell carcinoma arising from a native kidney with liver and bone metastasis in a renal transplant patient. World J Nucl Med 2018;17:296-8.

Access this article online	
<b>Website:</b> www.wjnm.org	<b>Quick Response Code</b> 
<b>DOI:</b> 10.4103/wjnm.WJNM_65_17	

## DISCUSSION


The incidence of RCC in native kidneys of renal transplant patients varies between 0.3% and 4.8%. The risk is increased in patients with acquired cystic kidney disease (ACKD), in men, recipients aged at least 65 years, those with a longer pretransplant dialysis interval, a donor aged at least 50 years, and microscopic hematuria.<sup>[2]</sup>

In patients with ACKD, the prevalence is 19% and in patients with complex cysts, it is 54%. Papillary hyperplasia of cyst epithelium is recorded in virtually every detailed pathology report of tumors arising in ACKD and is the likely pathogenetic basis for the development of renal tumors in cystic kidneys complicating dialysis.<sup>[3]</sup> A recent study showed that the occurrence of native renal cysts that is even a single cyst confers a 1.7-fold higher risk of developing RCC in renal transplant recipients.<sup>[4]</sup>

The influence of immunosuppressive agents on cyst formation remains poorly understood. Unlike other locations, immunosuppression appears not to increase the risk of developing malignant changes in native kidneys, as suggested by the identical incidence of RCC in heart transplant patients and the general population.<sup>[5]</sup>

The diagnosis of native kidney RCC in renal transplant recipients is difficult. It is typically an incidental finding during ultrasound scan or computed tomography for other clinical indications. In fact, most tumors are small and asymptomatic at presentation. In a retrospective study, the median native kidney tumor size was 2 cm.<sup>[6]</sup> Surgery is the preferred treatment as it is curative for the majority of RCC without metastasis. The patient showed liver and skeletal metastasis, so he was started on targeted therapy.

Unlike for most other malignancies, application of FDG PET/CT is limited for RCC, mainly due to physiological excretion of FDG from the kidneys, which decreases the contrast between renal lesions and normal tissue, and may obscure or mask the lesions of the kidneys. FDG PET/CT has potency as an imaging biomarker to provide useful information about patient's survival. Pretreatment SUV<sub>max</sub> assessed using FDG PET/CT can provide helpful information for clinical decision-making as it can serve as a useful prognostic marker for patients with advanced RCC. High SUV<sub>max</sub> in patients with primary RCC is suggested with correlate with a high likelihood of metastasis, and FDG accumulation may be useful in estimating patient's survival.<sup>[7]</sup> There are still no case reports of FDG PET/CT showing native kidney RCC in a transplant patient with liver and bone metastasis.



**Figure 1: Whole body Fluoroine-18 Fluorodeoxyglucose positron emission tomography/computer tomography showing maximum intensity projection image (a), coronal positron emission tomography/computer tomography (b) showing right native kidney primary renal cell carcinoma, axial positron emission tomography/computer tomography (c) showing liver lesion and axial positron emission tomography/computer tomography (d) showing bone metastasis (arrows)**

In conclusion, metastatic RCC can present as native kidney mass in a renal transplant recipient with liver and skeletal metastasis. F-18 FDG PET/CT scan allows detection of the metastatic sites making it a powerful diagnostic tool for an assessment of the extent of disease in patients of aggressive RCC.

### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

### REFERENCES

1. Kasiske BL, Snyder JJ, Gilbertson DT, Wang C. Cancer after kidney transplantation in the United States. *Am J Transplant* 2004;4:905-13.
2. Klatte T, Marberger M. Renal cell carcinoma of native kidneys in renal transplant patients. *Curr Opin Urol* 2011;21:376-9.

3. Schwarz A, Vatandaslar S, Merkel S, Haller H. Renal cell carcinoma in transplant recipients with acquired cystic kidney disease. *Clin J Am Soc Nephrol* 2007;2:750-6.
4. Goh A, Vathsala A. Native renal cysts and dialysis duration are risk factors for renal cell carcinoma in renal transplant recipients. *Am J Transplant* 2011;11:86-92.
5. Cheung CY, Lam MF, Lee KC, Chan GS, Chan KW, Chau KF, *et al.* Renal cell carcinoma of native kidney in Chinese renal transplant recipients: A report of 12 cases and a review of the literature. *Int Urol Nephrol* 2011;43:675-80.
6. Doublet JD, Peraldi MN, Gattegno B, Thibault P, Sraer JD. Renal cell carcinoma of native kidneys: Prospective study of 129 renal transplant patients. *J Urol* 1997;158:42-4.
7. Liu Y. The place of FDG PET/CT in renal cell carcinoma: Value and limitations. *Front Oncol* 2016;6:201.