

# Assessment of factors affecting the sleep hygiene of medical students in Bahawalpur, Pakistan: a cross-sectional study

Zarafshan Shafique\*  
Faiza Syed  
Safia Naz  
Saba Urooj  
Sadia Khan  
Sabahat Javed

Quaid-e-Azam Medical College,  
Department of Community Medicine -  
Bahawalpur - Punjab - Pakistan.

## ABSTRACT

**Objective:** To identify the factors which have a statistically significant association with sleep hygiene of medical students. **Material and Methods:** This study was done on 100 medical students. The questionnaire that was used for the collection of data had two parts. First was related to demographic variables and second was a validated Pittsburgh sleep quality index (PSQI). All participants were from the first to fifth years of Bachelor of Medicine and Bachelor of Surgery (MBBS). Data were analysed using SPSS 23. The association was found by applying chi-square test. **Results:** Our study revealed that out of 100 students, 61% had poor sleep hygiene (PSQI  $\geq$  5). Global PSQI scores among women were slightly higher than males. A statistical association was found between the residence and habit of studying at night of medical students with their sleep hygiene. **Discussion:** Medical students should ensure good sleep hygiene in order to maintain their academic performance, physical health, and mental health.

**Keywords:** Sleep Hygiene; Medical Students; Sleep Quality; Sleep Disturbance.

**\*Corresponding author:**  
Zarafshan Shafique  
E-mail: zarafshan.shafique23@gmail.com

Received: July 15, 2020;  
Accepted: October 5, 2020.

DOI: 10.5935/1984-0063.20200063

## INTRODUCTION

Sleep is defined as a natural state of rest during which the subject is unconscious but can be awoken with the help of proper external stimuli<sup>1</sup>. It is well known that a good quality and adequate amount of sleep are basic human necessities. According to sleep specialists, most healthy young adults and adults require at least seven to nine hours of sleep per day for optimal health and functioning of their mind and bodies<sup>2</sup>.

Sleep hygiene refers to those behaviours that ensure an improved quality and quantity of sleep to a person on a regular basis<sup>1</sup>. Reports show that about one-third of adults of the general population have some form of insomnia. Being a subgroup of the general population medical students are at an increased risk of developing a poor sleep hygiene, which, is mostly attributed to their challenging academic and clinical duties, long duration of study hours, stress of examination, hectic routine, and lifestyle choices<sup>3</sup>. Excessive use of caffeine, alcohol or energy drinks and unrestrained use of mobile phones, tablets or laptops before sleep are also some of those common factors that prevent maintaining good sleep hygiene<sup>4</sup>.

Most of the medical students do not give much importance to their sleep as they think that reducing the sleeping hours will help them cope with the piles of academic work that needs to be done in a short period of time. As a result, they develop poor sleeping habits in the weeks before the commencement of their examination and these altered sleeping patterns persist even after the examination is over<sup>5</sup>.

Studies have shown that deprivation of sleep results in impaired consciousness, sleepiness during daytime, lack of attention, compromised relationships, and fatigue<sup>6</sup>. These ill effects may in turn cause the students to commit medical errors and influence the patient safety<sup>7</sup>.

Sleep disturbances in medical students lead to such unwanted consequences that badly affect their academic performance, learning behaviours, and overall health<sup>8</sup>. Therefore, the importance of sleep quality among medical students cannot be overlooked as it has an obvious impact on the mental health, stress levels, and patient care<sup>9</sup>.

The objective of this study was to highlight those factors that have a statistically significant association with sleep hygiene. Since only a limited work has been done in Pakistan regarding this topic, our research will not only help doctors in expanding their knowledge but it will also serve to signify the importance of this matter in an attempt to improve the sleep hygiene of medical students across the country.

## MATERIAL AND METHODS

This cross-sectional study was conducted at Quaid-e-Azam Medical College, Bahawalpur after approval from the ethical review committee. A total of 100 medical students were recruited who were willing to participate in the study. The duration of study was from March 2020 to June 2020. The data were collected face-to-face using a self-administered questionnaire, which comprised of two parts (Questionnaire 1). Part one had seven questions related to demographic variables

and part two was a preformed pretested Pittsburgh sleep quality index (PSQI), which is a great tool for the evaluation of sleep hygiene during the past month. It consists of 19 items, which when combined yield seven sub-scales. Scoring of each item is done on a 3 point scale. The combination of these seven sub-scales produces a global PSQI score whose value ranges from 0 to 21. A global PSQI score of less than five indicated 'good' sleep hygiene in that subject while a score of equal to or greater than five meant 'bad' sleep hygiene<sup>10</sup>.

Data were analysed using SPSS v. 23. Frequencies and percentages were calculated for qualitative variables. The chi-square test was applied to test the significance of association between categorical variables. A *p*-value of less than 0.05 was considered as statistically significant.

The associated factors were exploited using both parts of the applied questionnaire. **Inclusion criteria:** all the students from first to fifth years of Bachelor of Medicine and Bachelor of Surgery (MBBS) who were willing and gave consent to participate in the study. **Exclusion criteria:** The students from first to fifth years of Bachelor of Medicine and Bachelor of Surgery (MBBS) who were not willing to participate and were not available during the study duration.

## RESULTS

In our study, there were 49 males and 51 females. Majority, i.e., 78% was resident of hostel (on-campus). Equal participants (20%) were chosen from each year of MBBS. The mean age of study participants was  $21.13 \pm 1.824$  years.

Out of 100 students, 61% were found to have poor sleep hygiene (their PSQI score was either equal to or more than five). Global PSQI scores among women were slightly higher than males (54.09% of females vs. 45.9% of males). Use of sleep medications was higher in males. Six male students vs. none of the female students used medication to sleep **less than once a week** (this is an idiomatic phrase and it was as such mentioned as one of the options to some questions in the PSQI questionnaire. It practically means that they **did not** use medication **every week**, rather once or twice a **month**). There was no difference in sleep efficiency of both genders.

### Components of Pittsburgh sleep quality index (PSQI)

#### Subjective sleep quality

According to Figure 1, 80% students had good sleep while 20% students regarded their sleep quality as bad.


#### Sleep latency

14% of study subjects it took to fall asleep within 15 minutes each night while 84% used to fall asleep after 15 minutes to more than an hour. 46% had trouble getting to sleep within 30 minutes less than once a week.

#### Sleep duration

30% used to spend more than seven hours in bed. 26% spent six to seven hours, 30% spent five to six hours while 14% spent less than five hours in bed.

**Rate of subjective sleep quality**



**Figure 1.** Rate of subjective sleep quality of medical students.

**Sleep efficiency**

71% of study subjects had sleep efficiency greater than 85%.

**Sleep disturbance**

23% used to wake up in the middle of the night once a week, which disturb their sleep, while 19% had disturbed sleep because of having bad dreams once or twice a week.

**Daytime dysfunction**

Out of 100 students, 41% had trouble staying awake less than once a week while driving, eating meals and engaging in social activities while it had been somewhat of a problem for 30% to keep up enough enthusiasm to get the things done.

**Use of sleep medications**

92% of the students did not use any type of sleep medication during the past month. 6% used sleeping medications less than once a week (**zero times a week but once or twice in the past month**) while only 2% used them once or twice a week.

As shown in Table 1, the p value obtained was 0.029 (which is less than 0.05), which meant that the association between sleep hygiene and the residence of medical students was significant. The students who resided on-campus were more likely to have bad sleep hygiene as compared to those who resided in their homes.

**Table 1.** Association of residence with sleep hygiene.

Residence	Global PSQI score		Total
	< 5	≥5	
On-campus	26	52	78
Off-campus	13	9	22
Total	39	61	100

( $\chi^2=4.786$ ; df-1;  $p=0.029$ ).

According to Table 2, the calculated p value was 0.020, which confirmed that those students who studied late at night actually had bad sleep hygiene as their global PSQI scores were comparatively greater than 5.

**Table 2.** Association of studying at night with sleep hygiene.

Studying at night	Global PSQI score		Total
	< 5	≥5	
Yes	26	52	78
No	13	9	22
Total	39	61	100

( $\chi^2=5.450$ ; df-1;  $p=0.020$ ).

Sleep hygiene of medical students did not seem to be affected by their gender as shown in Table 3.

**Table 3.** Association of gender with sleep hygiene.

Sleep hygiene	Male	Female	Total
Good sleep	21	18	39
Bad sleep	28	33	61
Total	49	51	100

( $\chi^2=0.601$ ; df-1;  $p=0.438$ ).

**DISCUSSION**

The findings of our study showed that 61% of all students had poor sleep hygiene. And 14% of students got less than 5 hours of sleep per night. In a study conducted by Waqas et al.<sup>11</sup>, on the association of academic stress with sleeping difficulties in medical students in Lahore (Pakistan), it was found that 77% of the respondents were poor sleepers. Academic stressors were a significant cause of poor sleep hygiene in these students. A high percentage, i.e., 27.8% of the total respondents used to get less than 5 hours of sleep per night<sup>11</sup>. An interesting finding of our study is that a statistical association was found between the residence and quality of sleep. The students who lived in hostels had comparatively bad sleep hygiene (85.24%) than those who lived in their homes (14.75%). This is opposite to the findings of a study conducted in West Indian Metropolitan city by Chutani et al.<sup>12</sup>, where more day scholars (73.7%) reported bad sleep quality as compared to hostellers (40.5%).

This could be due to the additional burden of travelling back and forth between college and home that consumes a lot of their time along with the academic load, which prevents them from getting a proper sleep. Hostellers, on the other hand, are provided with living accommodations in the vicinity of their college, so some of their time is saved that they can utilize in making up for their lost sleep due to extracurricular activities and odd sleeping hours<sup>12</sup>.

Our study reported that higher percentage of females (54.09%) had poor sleep hygiene as compared to males (45.9%). Our results were consistent with a study conducted in Dehradun which revealed that 59.3% of females had shown a poor quality of sleep, which was attributed to the long hours of mobile phone usage by them<sup>13</sup>.

No significant association was found between genders and sleep hygiene in our study. This is similar to the results of another study conducted on the evaluation of sleep hygiene among medical students in Saudi Medical College<sup>14</sup>.

In a study conducted by Alsaggaf et al.<sup>15</sup>, on sleep quantity, quality and insomnia symptoms of medical students

during clinical years in Kingdom of Saudi Arabia, it was reported that 30% of students had a poor sleep quality. Caffeinated beverages were taken by a high percentage (65%) of students during the day. This finding is consistent with our study where 54% of total students used to take coffee or tea every day (this was asked in first part of the applied Questionnaire 1).

A significant statistical association was found between the consumption of caffeinated drinks and poor sleep quality in a study conducted at King Abdul-Aziz University, Jeddah, KSA while no such statistical association was found in our study<sup>16</sup>.

Another finding of our study was that only a low percentage (6%) of students used to take sleep medications less than once a week and most of them were males. In a study conducted among Sudanese medical students by Mirghani et al.<sup>17</sup>, no difference was found for the use of sleep medications among excellent and average groups of students (23% among excellent group and 22.6% among average group).

There were some limitations in our study. The sample size was very small and it was done in only one institute so the results cannot be generalized. The effect of factors such as stress, anxiety and depression were not studied in this research. Therefore, we propose other researchers to take these points into consideration while conducting a study on sleep hygiene of medical students in future.

## CONCLUSION

In addition to the high prevalence of poor sleepers, our study has been able to find a significant association between the residence at hostel and habit of studying at night with the bad sleep hygiene of medical students. It is very important to arrange educational programs for the medical students in which they should be taught about the measures to improve their sleep hygiene and the ways through which they can fix their routines. Medical students should be assessed for their stress and anxiety profiles by their respective institutions and should be counselled accordingly.

## ACKNOWLEDGEMENT

I would like to acknowledge Dr. Faiza Syed for providing assistance in the collection of data, analysis of data

and proofreading the final results. She was extremely kind and helpful throughout the study.

## REFERENCES

1. Ananthachari KR, Sonavane R. Assessment of sleep hygiene among medical students at Gadag District, Karnataka. *Indian J Forensic Community Med.* 2017 Apr/Jun;4(2):77-80.
2. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health.* 2015 Mar;1(1):40-3.
3. Azad MC, Fraser K, Rumana N, Abdullah AF, Shahana N, Hanly PJ, et al. Sleep disturbances among medical students: a global perspective. *J Clin Sleep Med.* 2015 Jan;11(1):69-74.
4. Rezaei M, Khormali M, Akbarpour S, Hagighi KS, Shamsipour M. Sleep quality and its association with psychological distress and sleep hygiene: a cross sectional study among pre-clinical medical students. *Sleep Sci.* 2018 Jul/Aug;11(4):274-80.
5. Almojali AI, Almalki SA, Allothman AS, Masuadi EM, Alaqeel MK. The prevalence and association of stress with sleep quality among medical students. *J Epidemiol Glob Health.* 2017 Sep;7(3):169-74.
6. Lohitashwa R, Kadli N, Kisan R, Sindhuja A, Deshpande D. Effects of stress on sleep quality in young adult medical students: a cross sectional study. *Int J Res Med Sci.* 2015;3(12):3519- 23.
7. Siddiqui AF, Al-Musa H, Al-Amri H, Al-Qahtani A, Al-Shahrani M, Al-Qahtani M. Sleep patterns and predictors of poor sleep quality among medical students in King Khalid University, Saudi Arabia. *Malays J Med Sci.* 2016 Nov;23(6):94-102.
8. Subhaprada SC, Vijayakumari S, Rajasekhar T, Venkateswarlu U. An interventional study on sleep hygiene among medical students. *Int J Com Med Public Health.* 2017;4(2):352-56.
9. Arsalan B, Ahmed F. Sleep Hygiene and Quality among Medical Students. *IJR.* 2015;2(10):487-90.
10. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989 May;28(2):193-213.
11. Waqas A, Khan S, Sharif W, Khalid U, Ali A. Association of academic stress with sleeping difficulties in medical students of Pakistani medical school: a cross-sectional survey. *Peer J.* 2015 Mar;3:e840.
12. Chutani A, Shenvi DN, Singhal A. Sleep, sleepiness and medical college students: a comparative study among medical and paramedical students of a tertiary care teaching hospital from a West Indian metropolitan city. *Ann Med Health Sci Res.* 2017;7:85-91.
13. Yogesh S, Abha S, Priyanka S. Short communication, mobile usage and sleep patterns among medical students. *Indian J Physiol Pharmacol.* 2014 Jan/Mar;58(1):100-3.
14. Alqahtani JS, Al-Harbi R, Bathnain B, Al-Ghamdi F, Al-Shehri I, AlRabeeh S. D60. Sleep patterns in diverse populations. *Am J Respir Crit Care Med.* 2018;197:A7293.
15. Alsaggaf MA, Wali SO, Merdad RA, Merdad LA. Sleep quantity, quality, and insomnia symptoms of medical students during clinical years. *Saudi Med J.* 2016 Feb;37(2):173-82.
16. Ibrahim NK, Badawi FA, Mansouri YM, Ainousa AM, Jambi SK, Fatani AN, et al. Sleep quality among medical students at King Abdulaziz University: a cross-sectional study. *J Com Med Health Educ.* 2017;7(5):561.
17. Mirghani HO, Mohammed OS, Almurtadha VM, Ahmed MS. Good sleep quality is associated with better academic performance among Sudanese medical students. *BMC Res Notes.* 2015 nov;8:706.