

Sternoclavicular Joint Infection: Classification of Resection Defects and Reconstructive Algorithm

Janna Joethy¹, Chong Hee Lim², Heng Nung Koong³, Bien-Keem Tan¹

¹Department of Plastic, Reconstructive and Aesthetic Surgery, Singapore General Hospital; ²National Heart Centre; ³National Cancer Centre, Singapore

Background Aggressive treatment of sternoclavicular joint (SCJ) infection involves systemic antibiotics, surgical drainage and resection if indicated. The purpose of this paper is to describe a classification of post resectional SCJ defects and highlight our reconstructive algorithm. Defects were classified into A, where closure was possible often with the aid of topical negative pressure dressing; B, where parts of the manubrium, clavicular head, and first rib were excised; and C, where both clavicular, first ribs and most of the manubrium were resected.

Methods Twelve patients (age range, 42 to 72 years) over the last 8 years underwent reconstruction after SCJ infection. There was 1 case of a type A defect, 10 type B defects, and 1 type C defect. Reconstruction was performed using the pectoralis major flap in 6 cases (50%), the latissimus dorsi flap in 4 cases (33%), secondary closure in 1 case and; the latissimus and the rectus flap in 1 case.

Results All wounds healed uneventfully with no flap failure. Nine patients had good shoulder motion. Three patients with extensive clavicular resection had restricted shoulder abduction and were unable to abduct their arm past 90°. Internal and external rotation were not affected.

Conclusions We highlight our reconstructive algorithm which is summarised as follows: for an isolated type B SCJ defect we recommend the ipsilateral pectoralis major muscle for closure. For a type C bilateral defect, we suggest the latissimus dorsi flap. In cases of extensive infection where the thoracoacromial and internal mammary vessels are thrombosed, the pectoralis major and rectus abdominus cannot be used; and the latissimus dorsi flap is chosen.

Keywords Sternoclavicular joint / Infectious arthritis / Surgical flap

Correspondence: Bien-Keem Tan
Department of Plastic, Reconstructive and Aesthetic Surgery, Singapore General Hospital, Outram Road, Singapore 169608
Tel: +65 63214686
Fax: +65 62209340
E-mail: bienkeem@gmail.com

The authors thank Ms Jane Wong for the medical illustrations.

No potential conflict of interest relevant to this article was reported.

Received: 5 Jun 2012 • Revised: 10 Jul 2012 • Accepted: 17 Jul 2012

pISSN: 2234-6163 • eISSN: 2234-6171 • <http://dx.doi.org/10.5999/aps.2012.39.6.643> • Arch Plast Surg 2012;39:643-648

INTRODUCTION

Septic arthritis of the sternoclavicular joint (SCJ) is a rare clinical entity that represents 1% of all bone and joint infections [1,2]. Common predisposing factors include diabetes, rheumatoid arthritis, intravenous drug use, intraarticular injection, and immunosuppressive disorders [3]. It is often missed in patients with multiple comorbidities.

In early stages, the infection is sealed off by the joint capsule resulting in diagnostic delays. The inability of the capsule to distend leads to an initial build up of pressure within the joint which is subsequently released through fistulation, leading to cutaneous abscesses or mediastinitis [2,4]. Early diagnosis of SCJ infection requires a high index of suspicion and a confirmatory computed tomography (CT) scan.

Treatment is variable and ranges from simple drainage to ex-

tensive debridement as has been described in our earlier series [5,6]. We highlight our reconstructive approach to the management of cases where the debridement is substantial and the wound cannot be easily closed. In such cases, the defect size and exposure of critical structures dictate our reconstruction. To date, there is little in the literature detailing surgical reconstruction of SCJ infections. The purpose of this paper is to classify SCJ defects and describe our reconstructive algorithm.

METHODS

Twelve patients (7 female, 5 male) over the last 8 years underwent reconstruction after SCJ infection. The age ranged from 42 to 72 years old (mean, 59 years). A single case of bilateral SCJ infection is included and we discuss the reconstruction of this rare entity. After joint resection, negative pressure therapy was commenced to ensure optimization of patient and wound prior to reconstruction. The patients were optimized in terms of nutrition, sepsis and co-morbidities. Once both wound and patient were ready, reconstruction was performed according to our algorithm (Fig. 1).

RESULTS

Diabetes (n=7) followed by hypertension (n=4) were the commonest co-morbidities but often our patients had other medical conditions (Table 1). The commonest pathogen identi-

fied was *Staphylococcus aureus* (*S. aureus*, n=6) of which 2 were methicillin resistant. *Escherichia coli* (*E. coli*, n=3), *Streptococcus pyogenes* (*S. pyogenes*, n=2) and *Pseudomonas aeruginosa* (*P. aeruginosa*, n=1) were the others.

In these unwell patients, we started broad spectrum antibiotics




Table 1. Showing our patients, their co-morbidities, type of defect, the reconstruction and the functional outcome

Patient No.	Age (yr)/ Sex	Risk factors	Cause	Type of defect ^{a)}	Reconstruction	Outcome	Average follow-up (mo)
1	57/F	Systemic lupus erythematosus, interstitial lung disease, asthma	Central line insertion	A	Secondary closure	Uneventful recovery	24
2	65/F	Rheumatoid arthritis and a previous cerebrovascular accident	Sepsis	B	Latissimus dorsi flap	Uneventful recovery	36
3	45/M	Non insulin dependent diabetes mellitus	Sepsis	B	Pectoralis major flap	Uneventful recovery	36
4	46/M	Hypertension and hypercholesterolemia	Sepsis	B	Contralateral pectoralis major flap	Uneventful recovery	18
5	53/F	Non insulin dependent diabetes mellitus, hypothyroidism and hypertension	Unknown	C	Latissimus dorsi and rectus abdominis muscle flap	Uneventful recovery	36
6	62/F	Ischaemic heart disease	Central line insertion	B	Pectoralis major flap	Frozen shoulder	48
7	58/M	Hypertension, asthma	Central line insertion	B	Pectoralis major flap	Uneventful recovery	24
8	68/M	Insulin dependent diabetes mellitus	Sepsis	B	Latissimus dorsi muscle flap	Uneventful recovery	60
9	74/M	Renal failure, non insulin dependent diabetes mellitus	Sepsis	B	Pectoralis major flap	Decreased shoulder abduction	48
10	74/F	Insulin dependent diabetes mellitus	Sepsis	B	Pectoralis major flap	Decreased shoulder abduction	36
11	63/F	Renal failure	Central line insertion	B	Latissimus dorsi flap	Uneventful recovery	18
12	42/F	Non insulin dependent diabetes mellitus, hypothyroidism, hypertension	Sepsis	B	Latissimus dorsi flap	Uneventful recovery	18

^{a)}Type of defect refers to our classification of sternoclavicular defects (Fig. 5). In type A defects, the defect is small and closure is possible with the aid of topical negative pressure dressing. In type B defects, part of the manubrium, clavicular head and first rib are excised. In type C defects, the defect is bilateral involving both clavicular and first rib and most of the manubrium.

until sensitivities were available. High dose cloxacillin (*S. aureus*), vancomycin (methicillin resistant *S. aureus*) or penicillin (*S. pyogenes*) were used. Gram negative bacteria were treated with imipenem or tazocin (*P. aeruginosa*, *E. coli*). All antibiotic usage was used in consultation with infectious disease physicians.

Reconstruction was performed by using the pectoralis major flap in 6 cases (50%), the latissimus dorsi flap in 4 cases (33%), secondary closure in 1 case and another required two flaps: the latissimus dorsi and the rectus flap. All wounds healed uneventfully and there was no flap failure.

Three patients had restricted shoulder abduction and were unable to abduct their arm past 90 degrees. These patients had extensive clavicular resection. Internal and external rotation were not affected.

We highlight a few illustrative cases as follows.

Case 1

Type B defect (Fig. 2): latissimus dorsi musculocutaneous flap reconstruction. A 65-year-old female with rheumatoid arthritis and a previous cerebrovascular accident presented with sepsis. She underwent a CT chest which showed a collection at her

left SCJ with air pockets and mediastinal involvement. The collection was drained and washed out. Part of the manubrium, two-thirds of the clavicle and first rib were excised. As part of debridement, the left internal mammary artery and thoracoacromial artery were ligated. A latissimus dorsi flap was used to reconstruct this defect.

Case 2

Type B defect (Fig. 3): contralateral pectoralis major musculocutaneous flap reconstruction. A 46-year-old man with hypertension and hypercholesterolemia presented with multiple lymph nodes at the cervical chain and left axilla. A CT neck revealed a left SCJ infection with multiple abscesses involving the sternocleidomastoid muscles and the left pectoralis major muscle. He underwent excisional debridement and coverage using the contralateral pectoralis major.

Case 3

Type C defect (Fig. 4): latissimus dorsi and rectus abdominus muscle flap reconstruction. A 53-year-old lady with diabetes, hypothyroidism and hypertension presented with diffuse swelling






Fig. 4. Type C defect

Latissimus dorsi and rectus abdominus reconstruction. A 53-year-old lady with bilateral sternoclavicular joint (SCJ) infection. (A) Showing the wound following debridement with excision of the entire manubrium and upper half of the sternum. (B) After tunnelling the latissimus dorsi (LD) muscle on the left and rectus abdominus (RA) muscle on the right. (C) One year postoperative view.



over her chest. A CT confirmed bilateral SCJ infection. She underwent wound debridement with resection of her manubrium and upper sternum, bilateral clavicular heads and, both first ribs. The large residual defect was closed with a latissimus dorsi and rectus abdominus muscle flap.

DISCUSSION

Existing literature categorizes SCJ infection but not SCJ resection [8,9]. We have classified the post resectional SCJ defect and based our treatment accordingly. From our series, three types of defects exist (Fig. 5).

Type A


In early, well contained infections, the joint is stable and CT scanning shows infection confined to the joint with joint destruction and thickening of periarticular tissues. Following drainage and limited SCJ debridement (Figs. 5A, 6), the defects are small and lend themselves to secondary closure often with the aid of negative pressure dressing [2,7]. In such cases, we preserved the overlying pectoralis major for later coverage.

Type B

In extensive unilateral infection (Figs. 2, 7), there is spread beyond the joint to involve adjacent bone and rib as can be seen on CT with bony and joint destruction along with air pockets in subcutaneous layer or deeper in the anterior mediastinum. These cases will require at least excision of part of the clavicular head, part of the first rib, and part of the manubrium (Fig. 5B), leading to exposure of the thymus, brachiocephalic veins, and in some cases, the pleura. In these instances, we aim to cover the exposed bone ends along with any exposed underlying vital structures. We also introduce fresh blood supply in the form of a muscle flap to control infection. The preferred reconstructive

Fig. 5. Classification of sternoclavicular defects

In type A defects, closure is possible with the aid of topical negative pressure dressing. In type B defects, part of the manubrial head, clavicular head and first rib are excised. These defects are typically closed with an ipsilateral pectoralis major flap or an ipsilateral latissimus dorsi flap. In type C defects, the defect is bilateral involving both clavicular heads and first ribs and most of the manubrium. Closure here is performed with a latissimus dorsi flap and depending on the inferior extent of the wound, a rectus abdominis flap is used.




option was the ipsilateral pectoralis major flap based on the thoraco-acromial vessels. The pectoralis major was disinserted from the humerus and shifted medially to fill the defect. Externally the muscle was skin grafted if the wound was tight. In Fig. 3, where the infective process involved the ipsilateral pectoralis major, we used the contralateral pectoralis major with a long vertical skin paddle. The muscle alone could not reach the defect; it was the extended skin paddle that covered the wound. In cases where the thoraco-acromial vessels are of dubious viability, or when the vessels have been sacrificed during resection, we opted for latissimus dorsi flap reconstruction (Fig. 2).

Type C


More extensive infection resulted in resection of both clavicular

Fig. 6. Type A defect

No reconstruction. (A) A 57-year-old lady with systemic lupus erythematosus, interstitial lung disease and asthma had an internal jugular line inserted for postoperative monitoring. Three days post insertion, there was a tender swelling overlying her clavicle. (B) The strong anterior and posterior sternoclavicular ligaments wall off the infection and often result in a high pressure collection.

**Fig. 7. Type B defect**

Pectoralis major reconstruction. Showing a 45-year-old man with diabetes who presented with diabetic ketoacidosis secondary to sepsis. A computed tomography scan revealed a collection at the sternoclavicular joint. (A) Mobilisation of the pectoralis major to cover the defect. (B) Two months postoperative photo.



heads and both first ribs and the entire manubrium (Fig. 5C). The anterior chest became flail and there was loss of insertion of accessory muscles of ventilation. The priority here was to preserve the pectoralis major muscles to prevent destabilizing the chest and shoulders further. In Fig. 4, due to the large expanse of dead space, we used a latissimus dorsi and a rectus muscle. If the defect was confined to the manubrium, the latissimus dorsi flap alone would have been sufficient.

After SCJ resection, we constantly battle with instability of the shoulder girdle and movement of the thorax. This is in contrast with other regions like the lower extremity where immobilization is easily accomplished leading to earlier wound healing. There is little in the current literature for rigid fixation and joint replacement of the SCJ. The shoulder girdle articulates with the SCJ and with the removal of the medial clavicle, the clavicular

strut is lost affecting shoulder abduction. We noticed this in 3 patients, all of whom had long segment clavicular resection. One patient developed a frozen shoulder. Thus, we recommend aggressive physiotherapy 3 weeks after surgery.

Despite our elaborate reconstructive algorithm, the patients' comorbidities were taken into account. Most were critically ill with diabetes or chronic renal failure. In such cases, it was important to hold off surgery prior to optimization. Topical negative pressure dressing was used to bide time allowing improvement of nutrition, clearance of sepsis and control of comorbid conditions.

REFERENCES

1. Bar-Natan M, Salai M, Sidi Y, et al. Sternoclavicular infec-

- tious arthritis in previously healthy adults. *Semin Arthritis Rheum* 2002;32:189-95.
2. Ross JJ, Shamsuddin H. Sternoclavicular septic arthritis: review of 180 cases. *Medicine (Baltimore)* 2004;83:139-48.
 3. El Ibrahimy A, Daoudi A, Boujraf S, et al. Sternoclavicular septic arthritis in a previously healthy patient: a case report and review of the literature. *Int J Infect Dis* 2009;13:e119-21.
 4. Burkhart HM, Deschamps C, Allen MS, et al. Surgical management of sternoclavicular joint infections. *J Thorac Cardiovasc Surg* 2003;125:945-9.
 5. Lim CH, Lim J, Naik MJ, et al. Surgical management of sternoclavicular joint infection. *Asian Cardiovasc and Thorac Ann* 2001;9:291-5.
 6. Song HK, Guy TS, Kaiser LR, et al. Current presentation and optimal surgical management of sternoclavicular joint infections. *Ann Thorac Surg* 2002;73:427-31.
 7. Puri V, Meyers BF, Kreisel D, et al. Sternoclavicular joint infection: a comparison of two surgical approaches. *Ann Thorac Surg* 2011;91:257-61.
 8. Abu Arab W, Khadragui I, Echave V, et al. Surgical management of sternoclavicular joint infection. *Eur J Cardiothorac Surg* 2011;40:630-4.
 9. Nusselt T, Klinger HM, Freche S, et al. Surgical management of sternoclavicular septic arthritis. *Arch Orthop Trauma Surg* 2011;131:319-23.