LETTER

Multiple Sparganosis

Viroj Wiwanitkit

University Editorial Office, Hainan Medical University, Haikou, China

Correspondence: Viroj Wiwanitkit

University Editorial Office, Hainan Medical University, No. 3, College Road, Longhua District, Haikou, China

Tel: +86-6624132436, Fax: +86-6624132436, E-mail: wviroj@yahoo.com

No potential conflict of interest relevant to this article was reported.

Received: 17 Sep 2013 • Revised: 20 Oct 2013 • Accepted: 21 Oct 2013 pISSN: 2234-6163 • eISSN: 2234-6171 http://dx.doi.org/10.5999/aps.2014.41.2.181 • Arch Plast Surg 2014;41:181

Copyright © 2014 The Korean Society of Plastic and Reconstructive Surgeons This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dear Editor.

The recent report by Yoon et al. [1] on multiple sparganosis is very interesting. Generally, sparganosis is a parasitic infestation of tissue that is sporadically reported from many countries around the world. Yoon et al. [1] reported a case with this problem and concluded that "the longevity of spargana could have been up to 45 years with a parasitic infection occurring in two sites following a change in the patient's immune status". The main clinical presentation of sparganosis is a mass [2]. In some cases, it can be asymptomatic and is found by accident from pathohistological examination of a surgical specimen [2]. For management of sparganosis, complete surgical excision has been recommended [2]. Surgical treatment is usually effective and pathohistological examination of the surgical specimens is usually the appropriate way to obtain a final confirmatory diagnosis [2]. However, a chance of recurrence has been observed, and thus careful postoperative surveillance is recommended.

The longevity of spargana is worth noting. Up to the present, the exact lifespan of the parasite in humans has not been determined. In

the article in question, Yoon et al. [1] claimed that it was very strange that the parasite had survived for many years in their patient. The claim of long survival might be due to history taking, which could have been too subjective. Of interest, in a previous report by Wiwanitkit [2], only 14 of 34 patients could recall engaging in the risk behaviors (a history of drinking impure water, a history of eating frog or snake meat, or a history of using frog or snake meat as a poultice) [1]. It is very difficult to believe that the patient could remember a mundane event from 45 years earlier. In addition, a chronic parasitic infestation in a calcified lesion, such as a calcified tract, should be verified by pathohistological examination [3].

Another more interesting point is the existence of the disease in immunosuppressed patients. The effect of deteriorated immune status on this infection is remarkable. For another parasitic tissue infestation, cysticercosis, it was proposed that the outcome of infection might depend on the host's immunological status [4]. However, to the best of our knowledge, no report has been published of sparganosis in an human immunodeficiency virus-infected patient. In fact, multiple sparganosis can be seen in patients without impaired immune status [5]. The interrelationship between impaired immune status and sparganosis should thus be further investigated.

References

- 1. Yoon HS, Jeon BJ, Park BY. Multiple sparganosis in an immunosuppressed patient. Arch Plast Surg 2013;40:479-81.
- Wiwanitkit V. A review of human sparganosis in Thailand. Int J Infect Dis 2005;9:312-6.
- Lee KJ, Myung NH, Park HW. A case of sparganosis in the leg. Korean J Parasitol 2010;48:309-12.
- Hoberg EP, Ebinger W, Render JA. Fatal cysticercosis by Taenia crassiceps (Cyclophyllidea: Taeniidae) in a presumed immunocompromised canine host. J Parasitol 1999;85:1174-8.
- Moulinier R, Martinez E, Torres J, et al. Human proliferative sparganosis in Venezuela: report of a case. Am J Trop Med Hyg 1982;31:358-63.