

IMAGES

Late-Onset Methicillin-Resistant *Staphylococcus aureus* Infection after Facial Poly-L-Lactic Acid Injection

Min Choi, Ji Seon Cheon, Woo Young Choi,
Kyung Min Son

Department of Plastic and Reconstructive Surgery, Chosun
University College of Medicine, Gwangju, Korea

Correspondence: Ji Seon Cheon
Department of Plastic and Reconstructive Surgery, Chosun University College
of Medicine, 309 Pilmun-daero, Dong-gu, Gwangju 61452, Korea
Tel: +82-62-220-3180, Fax: +82-62-225-0996, E-mail: ps9107@naver.com

No potential conflict of interest relevant to this article was reported.

Received: 7 Dec 2016 • Revised: 16 Feb 2017 • Accepted: 4 Mar 2017
pISSN: 2234-6163 • eISSN: 2234-6171
<https://doi.org/10.5999/aps.2017.44.3.248>
Arch Plast Surg 2017;44:248-249



Copyright © 2017 The Korean Society of Plastic and Reconstructive Surgeons
This is an Open Access article distributed under the terms of the Creative Commons
Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial use, distribution, and reproduction in any
medium, provided the original work is properly cited.

Poly-L-lactic acid (PLLA) is widely used in lipotrophy and facial rejuvenation for its ability to stimulate collagen neogenesis. Although its safety and efficacy have been established, adverse effects have been reported [1,2]. Late-onset adverse effects, occurring 6 months or more after injection, include granulomas, nodules, and pseudoabscess [1].

A 43-year-old woman presented with painful erythematous nodules on the left temple (Fig. 1), where PLLA had been injected for augmentation 14 months previously. She had no predisposing factors, and empirical antibiotic treatment did not lead to improvement. After incision of the nodules, pus and tissue materials were drained (Fig. 2), Histopathologic findings showed an abscess without granulomatous features around foreign body

materials (Fig. 3), and methicillin-resistant *Staphylococcus aureus* (MRSA) was cultured. We therefore made the diagnosis of late-onset MRSA infection after PLLA injection.

In this case, the subcutaneous nodules showed no sign of a granulomatous reaction, and MRSA was grown in a bacterial culture. Bacterial contamination of the filler materials may have been the cause of these nodules, which is related to the biofilm theory. Biofilms, defined as communities of microorganisms, are attached to the inert surfaces of catheters, and *Staphylococcus aureus* is frequently found in a dormant state within biofilms. It can be reactivated by external triggering factors (e.g., trauma), causing low-grade infection and abscess [3]. Our patient had not undergone any additional procedures, but repetitive traumas in daily life may have served as the triggering factor. Intravenous vancomycin was administered accordingly, and the



Fig. 2. After incision of the nodule, material including pus and tissue was drained.




Fig. 3. Skin biopsy revealing numerous inflammatory cells with fragments of foreign materials (H&E, ×200; neutrophil infiltration [red arrowhead], foreign body [blue arrow]).




Fig. 1. Several painful erythematous nodules (red arrowhead) on the left temporal area of the patient's face.

nodules recovered completely without scarring. Therefore, physicians who perform aesthetic procedures should be aware that late-onset painful nodules are likely to be caused by a bacterial infection such as MRSA, and identification of the bacteria may be preferable to hastily pursuing other treatment possibilities.

References

1. Alijotas-Reig J, Garcia-Gimenez V, Vilardell-Tarres M. Late-onset immune-mediated adverse effects after poly-L-lactic acid injection in non-HIV patients: clinical findings and long-term follow-up. *Dermatology* 2009;219:303-8.
2. Choi WY, Cho HW, Lee DW. Complications of injectable soft tissue filler. *Arch Aesthetic Plast Surg* 2015;21:1-6.
3. Sadashivaiah AB, Mysore V. Biofilms: their role in dermal fillers. *J Cutan Aesthet Surg* 2010;3:20-2.

spontaneous pseudoaneurysm of the proximal occipital artery that presented as a neck mass.

A 44-year-old male patient presented with a pulsatile swelling lesion in the neck that had grown spontaneously over the course of a year (Fig. 1). The swelling had gradually progressed, and he had no previous history of head trauma, infection, or a genetic disorder. The patient was found to have a firm, non-tender, pulsating mass in the left side of the neck that measured 4 × 3 cm. Ultrasonography showed turbulent flow and dilatation of the involved vessel. Neck computed tomography showed partial opacification of the swelling and revealed a pseudoaneurysm arising from the left proximal occipital artery (Fig. 2). The decision was made to embolize the lesion via angiography (Fig. 3). The mass was treated successfully by endovascular coil embolization (Fig. 4). One year postoperatively, the swelling had improved and recurrence was not detected.

Most occipital artery pseudoaneurysms are consequences of trauma, or are secondary to

Spontaneous Pseudoaneurysm of the Proximal Occipital Artery Presenting as a Neck Mass

Tae Gon Kim¹, Jin Ho Lee¹, Kyu Jin Chung¹, Jun-Ho Lee¹, Yong-Ha Kim¹, Chul Hoon Chang²

Departments of ¹Plastic and Reconstructive Surgery and ²Neurosurgery, Yeungnam University College of Medicine, Daegu, Korea

Correspondence: Tae Gon Kim
Department of Plastic and Reconstructive Surgery, Yeungnam University College of Medicine, 170 Hyeonchung-ro, Nam-gu, Daegu 42415, Korea
Tel: +82-53-620-3483, Fax: +82-53-626-0705
E-mail: kimtg0919@hanmail.net

No potential conflict of interest relevant to this article was reported.

Received: 23 Feb 2017 • Revised: 28 Mar 2017 • Accepted: 28 Mar 2017
pISSN: 2234-6163 • eISSN: 2234-6171
<https://doi.org/10.5999/aps.2017.44.3.249>
Arch Plast Surg 2017;44:249-250

Copyright © 2017 The Korean Society of Plastic and Reconstructive Surgeons
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Occipital artery pseudoaneurysms are rare vascular lesions, and are most often reported as a consequence of trauma [1,2]. We report the rare case of a



Fig. 1. Preoperative photograph showing a lesion on the left neck.




Fig. 2. Computed tomography scan (axial section) shows a left proximal occipital artery pseudoaneurysm.