COMMUNICATIONS

A Novel Method for Intraoperative Breast Implant Pocket Assessment: Air Augmentation

Dominik Duscher¹, Raphael Wenny¹, Francesco Somma², Georg M. Huemer¹

¹Section of Plastic, Aesthetic and Reconstructive Surgery, Kepler University Hospital Linz, Linz, Austria; ²Private Practice, Sorrento, Italy

Correspondence: Georg M. Huemer

Section of Plastic, Aesthetic and Reconstructive Surgery, Kepler University Hospital Linz, Krankenhausstrasse 9, A-4020 Linz, Austria Tel: +43-732-7806-73819, Fax: +43-732-7806-3190, E-mail: ghuemer@drhuemer.com

This article contains Supplemental Video S1.

No potential conflict of interest relevant to this article was reported.

Received: 28 Dec 2016 • Revised: 5 Feb 2017 • Accepted: 6 Feb 2017 pISSN: 2234-6163 • eISSN: 2234-6171 https://doi.org/10.5999/aps.2017.44.4.354 • Arch Plast Surg 2017;44:354-355

Copyright © 2017 The Korean Society of Plastic and Reconstructive Surgeons This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dear Editor-in-Chief,

The estimation of breast implant pocket size in aesthetic breast surgery is often a matter of clinical and intraoperative trial and error, which can negatively influence postoperative outcomes. While there are numerous preoperative and intraoperative techniques and devices to determine the ideal implant volume [1], it can be challenging to evaluate pocket preparation accurately. Herein, we describe a new method of elegantly assessing the extent and quality of implant pocket dissection intraoperatively.

An ideal implant pocket is defined by having a size adequate to fit the chosen prosthesis snugly without being too small, resulting in a distorted implant shape, or being too big, potentially leading to dislocation [2]. To determine if the implant pocket has been dissected sufficiently, we developed a novel technique based on temporally trapping air within the pocket. Employing this air augmentation method allows the progress of pocket preparation to be assessed accurately without substantial technical effort, and provides the opportunity for modifications based on the desired implant volume. Furthermore, it enables the surgeon to judge the edges of the implant pocket, making it a fast and inexpensive technique for assessing the symmetry and extent of the pocket shape.

Using the air augmentation technique, the surgeon is able to check for potential instances of adherence, unevenly dissected edges, and boxiness of the implant pocket before moving forward with breast implant placement. By simply inserting the index and middle finger into the inframammary wound, lifting up his or her hand, and quickly lowering it again air can be trapped into the dissected cavity via a vacuum effect (Fig. 1). The size and shape of the inflated pocket can be evaluated easily (Supplemental Video S1). This is especially helpful in judging the extent of medial release of the muscle and the boxiness of the lateral extent of the implant pocket. The inflated air

Fig. 1. Pocket size estimation using the air augmentation technique

(A, B) A 28-year-old patient underwent aesthetic breast surgery. After dissection of the implant pocket, the surgeon's index and middle finger were inserted into the inframammary wound. (C) The hand was lifted to allow the influx of air into the implant pocket. (D) Quickly lowering the hand while leaving the fingers in the pocket, acting as a plug, trapped the air within the pocket, enabling assessment of the dissected area.



can be forced against the boundaries of the implant cavity, mimicking the shape of the implant.

The air augmentation technique may help with dissection of the pocket, but it is not itself sufficient to support final decisionmaking about the implant size. However, in our practice, implant size is determined via a thorough preoperative investigation of the patient's body type and aesthetic goals. Relying on the preoperatively determined implant size combined with a precise intraoperative assessment of the adequacy of the pocket size and shape using the air augmentation technique has enabled us to completely omit traditional approaches using trial sizers, which are expensive and timeconsuming, and potentially have adverse effects such as increased complication rates and wound edge injury due to friction forces [3,4]. The air augmentation method requires only a short period of practice and is easily reproducible, making it a useful addition to the technical armamentarium of both novice and expert surgeons.

References

- 1. Caulfield RH, Niranjan NS. Innovative techniques: a novel technique for intraoperative estimation of breast implant size in aesthetic and reconstructive breast surgery. Aesthetic Plast Surg 2008;32:126-9.
- Hidalgo DA, Spector JA. Breast augmentation. Plast Reconstr Surg 2014;133:567e-583e.
- 3. Khoo LS, Radwanski HN, Senna-Fernandes V, et al. Does the use of intraoperative breast sizers increase complication rates in primary breast augmentation? A retrospective analysis of 416 consecutive cases in a single institution. Plast Surg Int 2016;2016:6584810.
- Holmes WJ, Timmons MJ, Kauser S. Techniques used by United Kingdom consultant plastic surgeons to select implant size for primary breast augmentation. J Plast Reconstr Aesthet Surg 2015;68:1364-9.

Supplemental Video S1. Air augmentation performed intraoperatively to assess implant pocket dissection.

Supplemental data can be found at: http://e-aps.org/src/sm/aps-44-354-s001.mp4